

AD-A210 118

REPORT DOCUMENTATION PAGE

REPORT DOCUMENTATION PAGE					
1. OFFICIAL USE ONLY.	2. REPORT DATE.	3. TYPE OF REPORT AND DATES COVERED.			
	November 1988	Final			
Significant Digit Computation of the Incomplete Beta Function Ratios					
Armido R. DiDonato Alfred H. Morris, Jr.					
Naval Surface Warfare Center (K33) Dahlgren, VA 22448-5000		NSWC TR 88-365			
Approved for public release; distribution is unlimited.					
<p style="text-align: right;">Sub X / Sub X /</p> <p>An algorithm is given for evaluating the incomplete beta function ratio $I_x(a,b)$ and its complement $1-I_x(a,b)$. Two new procedures are used with classical results. A listing of a transportable Fortran subroutine using this algorithm is given. The subroutine is accurate to 14 significant digits when the precision is not restricted by inherent error.</p> <p style="text-align: right;"><i>Keywords: incomplete beta function ratio, Fortran software</i></p>					
incomplete beta function ratio subroutine Fortran software			82		
UNCLASSIFIED	UNCLASSIFIED	UNCLASSIFIED			

FOREWORD

The work described in this report was done in the Space and Surface Systems Division and the Computer and Information Systems Division of the Strategic Systems Department. Its purpose was to design Fortran software, suitable for a high quality mathematics and/or statistics subroutine library, which yields the incomplete beta function ratios over an extensive range of the variables.

The authors are indebted to Dr. James Perry of NSWC for proving a result which is used in this work. His proof is given in Appendix A. The authors also wish to thank Ms. Dottie Burgess for making Figures 1 and 3.

Approved by:

R.L. Schmidt
R. L. SCHMIDT, Head
Strategic Systems Department



Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/ _____	
Availability Codes	
Dist	Avail and/or Special
A-1	

CONTENTS

	Page
I. Introduction.....	1
II. Basic Relations	3
III. Domains for Subprograms.....	11
IV. Auxiliary Functions	19
V. Concluding Remarks.....	23
VI Numerical Examples	25
References.....	27
 APPENDICES	
A. Bounds on $I_x(a,b)$ when $x=a/(a+b)$, $\min(a,b) \geq 1$	A-1
B. Monotonic Properties of $I_x(a,b)$	B-1
C. Uniform Asymptotic Behavior of (16)	C-1
D. Fortran Listing of BRATIO	D-1
DISTRIBUTION.....	(1)

I. INTRODUCTION

The incomplete beta function $I_x(a,b)$ is defined by

$$I_x(a,b) = G(a,b) \int_0^x t^{a-1} (1-t)^{b-1} dt, \quad a>0, b>0, 0 \leq x \leq 1 \quad (1)$$

$$B(a,b) = 1/G(a,b) = \int_0^1 t^{a-1} (1-t)^{b-1} dt = \Gamma(a)\Gamma(b)/\Gamma(a+b), \quad [10, p.36], \quad (2)$$

where the gamma function $\Gamma(u)$ is given by

$$\Gamma(u) = \int_0^\infty e^{-t} t^{u-1} dt, \quad u > 0. \quad (3)$$

Thus $I_1(a,b)=1$. In addition, if $0 < x < 1$ then $I_x(0,b)=1$ and $I_x(a,0)=0$.

The quantity $1 - I_x(a,b)$ is called the *complement* of $I_x(a,b)$. Using $u=1-t$ in (1) and (2) yields

$$1 - I_x(a,b) = I_y(b,a), \quad y = 1 - x. \quad (4)$$

The function I_x occurs in many branches of science, including atomic physics, fluid dynamics, transmission theory, lattice theory, and operations research. It is perhaps best known for its extensive applications in statistics. In particular, the well-known central F-distribution $P(F_0|\nu_1, \nu_2)$ can be obtained from I_x by the following substitutions :

$$a = \nu_1/2, \quad b = \nu_2/2, \quad t = \nu_1 F / [\nu_2 + \nu_1 F], \quad F_0 = \frac{bx}{a(1-x)}, \quad (5)$$

where

$$P(F_0|\nu_1, \nu_2) = \nu_1^{\nu_1/2} \nu_2^{\nu_2/2} G\left(\frac{\nu_1}{2}, \frac{\nu_2}{2}\right) \int_0^{F_0} F^{(\nu_1-2)/2} (\nu_2 + \nu_1 F)^{-(\nu_1+\nu_2)/2} dF \quad (6)$$

$$Q(F_0|\nu_1, \nu_2) = 1 - P(F_0|\nu_1, \nu_2) = I_{1-x}(\nu_2/2, \nu_1/2). \quad (7)$$

The incomplete beta function is also directly related to the Student's t-distribution $A(t_0|\nu)$ and the binomial distribution $E(n,r,x)$, where

$$P(|t| \leq t_0) = A(t_0|\nu) = \frac{2}{\sqrt{\nu}} G\left(\frac{1}{2}, \frac{\nu}{2}\right) \int_0^{t_0} (1+t^2/\nu)^{-(\nu+1)/2} dt \quad (8)$$

$$= 1 - I_x(\nu/2, 1/2), \quad x = \nu / [\nu + t_0^2],$$

$$E(n,r,x) = \sum_{i=r}^n \binom{n}{i} x^i (1-x)^{n-i} = I_x(r, n-r+1). \quad (9)$$

Derivations of these well-known results are given in [4].

Procedures for computing I_x date back to Newton. A historical survey outlining some of the analytical and approximation methods used for evaluating I_x is given by Dutka [7]. An extensive literature search for a robust algorithm to compute I_x did not reveal a publication that would lead to a subroutine acceptable for inclusion in a high quality main frame mathematics library such as the NSWC Library of Mathematics Subroutines (NSWCLIB), [9]. The best procedure found was in a report written by Amos and Daniel [2]. For a special case however, where a and b take positive integer and half-integer values, an algorithm exists, with an associated Fortran subroutine ISUBX, which yields I_x with 9-10 decimal-digit accuracy, [4, 5]. It is contained in NSWCLIB.

In this report an algorithm is given for computing I_x and $1 - I_x$. A transportable Fortran subroutine named BRATIO has been written which uses the algorithm. BRATIO is designed for use on computers having k -digit single precision floating arithmetics where $6 \leq k \leq 14$. On the CDC 6000-7000 series computers, BRATIO yields results accurate up to 14 significant digits for both I_x and $1 - I_x$. BRATIO is available for general use in the NSWC mathematics subroutine library, [9].

A primary region of difficulty for computing $I_x(a,b)$ and $1 - I_x(a,b)$ has been when a and/or b is large and $x \approx a/(a+b)$. In this region I_x changes rapidly from 0 to 1. A continued fraction, with new weighting factors, and a new asymptotic expansion are used to treat this region satisfactorily.

Section II contains the basic equations and algorithms used for BRATIO. Section III describes in a, b, x space the regions of use for these basic relations. An associated flowchart for BRATIO is included. Section IV describes a number of specialized algorithms required in order to use the basic relations effectively. Section V briefly summarizes the accuracy and efficiency of BRATIO, and section VI contains a few examples using BRATIO. Appendices A, B, and C contain proofs for some results which are used in BRATIO. A Fortran listing of BRATIO and its required subprograms is given in Appendix D.

II. BASIC RELATIONS

Throughout we shall use

$$p = a/(a+b) \quad q = 1 - p = b/(a+b). \quad (10)$$

Also since I_x and $1 - I_x$ are to be computed to the greatest possible accuracy,

$$y = 1 - x \quad (11)$$

is required as input in addition to a , b , and x . In BRATIO, relations (12)–(16) are used to compute either I_x or its complement. Their domains of application are given in Section III.

BPSER

$$I_x(a,b) \approx G(a,b) \frac{x^a}{a} \left(1 + a \sum_{j=1}^N \frac{(1-b)(2-b) \cdots (j-b)}{j! (a+j)} x^j \right) \quad (12)$$

The series is obtained from (1) by replacing the second factor in the integrand with its binomial expansion. Relation (12) is used only when $x \leq 0.7$ and $b \leq 1$, or $bx \leq 0.7$.

Given a tolerance $\epsilon > 0$, (12) is computed by the function BPSER(a,b,x,ϵ), where

$$I_x(a,b) \approx G(a,b) \frac{x^a}{a} \left(1 + a \sum_{n=1}^N w_n \right) \quad (12.1)$$

$$w_n = C_n / (a+n), \quad C_n = C_{n-1} (1-b/n)x, \quad C_0 = 1 \quad (12.2)$$

$N =$ the smallest integer such that $a|w_N| \leq \epsilon$.

BUP

$$I_x(a,b) - I_x(a+N,b) = x^a y^b \sum_{j=1}^N \frac{\Gamma(a+b+j-1)}{\Gamma(b)\Gamma(a+j)} x^{j-1}, \quad (N \geq 1) \quad (13)$$

Given a tolerance ϵ , (13) is computed by the function BUP(a,b,x,y,N,ϵ). BUP is used only with BPSER or the next relation to be described, BGRAT. Relation (13) follows from $I_x(a+1,b) = I_x(a,b) - x^a y^b G(a,b)/a$, which can be obtained by substituting

$$(a+b)t^a(1-t)^{b-1} = at^{a-1}(1-t)^{b-1} - \frac{d}{dt} [t^a(1-t)^b] \quad (13.1)$$

$$I_x(a+1,b) = \frac{a+b}{aB(a,b)} \int_0^x t^a(1-t)^{b-1} dt.$$

Equation (13) can be rewritten in the form

$$I_x(a,b) - I_x(a+N,b) = \frac{x^a y^b}{a B(a,b)} \sum_{i=0}^{N-1} d_i x^i, \quad d_{i+1} = \frac{a+b+i}{a+1+i} d_i, \quad d_0 = 1. \quad (13.2)$$

If $b \leq 1$ then for $i \geq 0$, $d_{i+1} \leq d_i$ and the sequence $h_i \equiv d_i x^i$ is monotonically decreasing. Thus, the computation of the sum $\sum h_i$ can be terminated when a term $h_m = d_m x^m$ is reached that satisfies $h_m \leq \epsilon \sum_{i=0}^m h_i$, or $m=N-1$.

When $b > 1$ then $h_i \geq h_{i+1}$ if and only if $x \leq (a+1+i)/(a+b+i) \equiv r_i$. Also we note that $r_i < r_j$ when $i < j$. Thus if k is the largest integer such that $x \geq r_{k-1}$, then

$$h_0 \leq h_1 \leq \dots \leq h_k \geq h_{k+1} \geq \dots \geq h_{N-1}.$$

Therefore, if k is the largest integer for which $k \leq (b-1)x/y - a$, then the computation of the sum $\sum h_i$ can be terminated when a term h_m ($m > k$) is met that satisfies $h_m \leq \epsilon \sum_{i=0}^m h_i$, or $m=N-1$.

BGRAT

$$I_X(a,b) \approx M \sum_{n=0}^L p_n J_n(b,u), \quad M = \frac{\Gamma(a+b)}{\Gamma(a)T^b} r, \quad a > b \quad (14)$$

$$T = a + \frac{b-1}{2}, \quad u = -T \ln x, \quad r = e^{-u} u^b / \Gamma(b), \quad J_0(b,u) = Q(b,u)/r \quad (14.1)$$

$$p_n = (b-1)c_n + \frac{1}{n} \sum_{m=1}^{n-1} (mb-n)c_m p_{n-m}, \quad c_m = \frac{1}{(2m+1)!}, \quad p_0 = 1 \quad (14.2)$$

$$Q(b,u) = \int_u^\infty \frac{e^{-t} t^{b-1}}{\Gamma(b)} dt \quad (\text{incomplete gamma function}) \quad [1; p.260] \quad (14.3)$$

$$J_{n+1}(b,u) = \frac{(b+2n)(b+2n+1)}{4T^2} J_n(b,u) + \frac{u+b+2n+1}{4T^2} \left(\frac{\ln x}{2}\right)^{2n} \quad (14.4)$$

$$L = \text{smallest integer such that } |p_L J_L| \leq \epsilon \left(\sum_{n=0}^L p_n J_n + w_0/M \right). \quad (14.5)$$

Relation (14) is used only when $a \geq 15$, $b \leq 1$, and $x > 0.7$. Given ϵ , (14) is computed by the subroutine BGRAT(a,b,x,y,w,ϵ , IERR) where w and IERR are variables. Given an initial value w_0 for w , then BGRAT assigns w the value $w_0 + I_X(a,b)$. IERR is an indicator which is set when underflow forces the computation of (14) to end prematurely.

Equation (14) was derived by Wise [15]. Our derivation follows:

If $t = e^{-y/T}$, then

$$\int_0^x t^{a-1} (1-t)^{b-1} dt = \frac{2^{b-1}}{T} \int_u^\infty e^{-y} \sinh^{b-1} \left(\frac{y}{2T}\right) dy \quad (14.6)$$

where T and u are defined above. Substituting

$$\sinh^{b-1} z = z^{b-1} \left(\sum_{n=0}^{\infty} c_n z^{2n} \right)^{b-1} = \sum_{j=n}^{\infty} p_j z^{2j+b-1} \quad (14.7)$$

into (14.6), where the c_n and p_j satisfy (14.2), yields (14) where

$$J_n(b, u) = \frac{1}{(2T)^{2n}} \frac{\Gamma(b+2n)}{r\Gamma(b)} Q(b+2n, u). \quad (14.8)$$

Then (14.4) follows from (14.8) and

$$Q(b+2n+2, u) = Q(b+2n, u) + \frac{e^{-u} u^{b+2n}}{\Gamma(b+2n+2)} (u+b+2n+1), \quad [1; 6.5.21] \quad (14.9)$$

where (14.9) is the result of two integrations by parts of its left hand side.

BFRAC

$$I_x(a, b) \approx \frac{x^a y^b}{B(a, b)} \left(\frac{\alpha_1}{\beta_1 +} \frac{\alpha_2}{\beta_2 +} \dots \frac{\alpha_m}{\beta_m} \right), \quad x \leq p \equiv \frac{a}{a+b} \quad (15)$$

$$\alpha_1 = 1, \quad \beta_1 = \frac{a}{a+1}(\lambda+1), \quad \lambda = a - (a+b)x = (a+b)(p-x) \quad (15.1)$$

$$\alpha_{n+1} = \frac{(a+n-1)(a+b+n-1)}{(a+2n-1)^2} n(b-n)x^2, \quad n \geq 1 \quad (15.2)$$

$$\beta_{n+1} = n + \frac{n(b-n)x}{a+2n-1} + \frac{a+n}{a+2n+1} [\lambda + 1 + n(1+y)], \quad n \geq 0 \quad (15.3)$$

$$\begin{cases} A_{n+1} = \beta_{n+1} A_n + \alpha_{n+1} A_{n-1} & A_0 = 0, \\ B_{n+1} = \beta_{n+1} B_n + \alpha_{n+1} B_{n-1} & B_0 = 1, \end{cases} \quad B_1 = \beta_1 \quad (15.4)$$

$$\frac{A_n}{B_n} = \frac{\alpha_1}{\beta_1 +} \frac{\alpha_2}{\beta_2 +} \dots \frac{\alpha_n}{\beta_n}. \quad (15.5)$$

m = the smallest integer such that

$$|A_m/B_m - A_{m-1}/B_{m-1}| \leq \epsilon |A_m/B_m|.$$

Equation (15) is new and is used over a very large part of the domain: $a > 1$, $b \geq 40$, with $x \leq p$. Given a tolerance ϵ , (15) is computed by the function BFRAC($a, b, x, y, \lambda, \epsilon$). The weighting factors c_n , given below, which control overflow and underflow, appear to be new.

Relation (15) is obtained by considering the classical expansion

$$I_X(a,b) = \frac{x^a y^b}{aB(a,b)} \left(\frac{1}{1} \frac{d_1}{1+1} \frac{d_2}{1+1+1} \dots \right) \quad (15.6)$$

$$d_{2n} = \frac{n(b-n)}{(a+2n-1)(a+2n)} x, \quad n > 0 \quad (15.7)$$

$$d_{2n+1} = -\frac{(a+n)(a+b+n)}{(a+2n)(a+2n+1)} x, \quad n \geq 0. \quad [1; 26.5.8], [3], [13] \quad (15.8)$$

For large a where $a \gg b$, we note that $d_{2n} \approx 0$ and $d_{2n+1} \approx -(1+b/a)x$. Thus if R_n is the iterate

$$R_n = \frac{1}{1} \frac{d_1}{1+1} \frac{d_2}{1+1+1} \dots \frac{d_{n-2}}{1+d_{n-1}},$$

then most of the change of values of these iterates occurs in every other iterate. Consequently, the "associated" continued fraction $\frac{a_1}{b_1} \frac{a_2}{b_2} \dots$ is considered where

$$\begin{aligned} a_1 &= 1, & a_{n+1} &= -d_{2n-1} d_{2n} \\ b_1 &= 1+d_1, & b_{n+1} &= 1+d_{2n}+d_{2n+1}, \quad n \geq 1. \end{aligned} \quad [14; p.20]$$

The iterates R_{2n} are the iterates for this expansion as already pointed out by Aroian in [3] (with some corrections given in [13]). Hence

$$I_X(a,b) = \frac{x^a y^b}{aB(a,b)} \left(\frac{a_1}{b_1} \frac{a_2}{b_2} \dots \right).$$

Now for large a where $a \gg b$, it is clear that $d_{2n} \approx 0$ forces $a_n \approx 0$. However, for $x \approx p$ we also find that $d_{2n} \approx 0$ forces $b_n \approx 0$. This can cause division of 0 by 0 when the iterates of this continued fraction are computed, or it can cause division to overflow. Thus, this continued fraction is also not considered appropriate for computational purposes.

In order to eliminate the problems arising from $d_{2n} \approx 0$, we rescale the coefficients a_n and b_n with weighting factors c_n such that

$$\bar{\alpha}_1 = c_1 a_1, \quad \alpha_n = c_{n-1} c_n a_n \quad (n \geq 2), \quad \beta_n = c_n b_n \quad (n \geq 1),$$

where

$$c_n = a + 2(n-1), \quad n \geq 1. \quad (15.9)$$

Then the iterates of $\frac{\bar{\alpha}_1}{\beta_1} \frac{\alpha_2}{\beta_2} \dots$ are the scaled iterates for $\frac{a_1}{b_1} \frac{a_2}{b_2} \dots$ and we obtain

$$I_X(a,b) = \frac{x^a y^b}{aB(a,b)} \left(\frac{\bar{\alpha}_1}{\beta_1} \frac{\alpha_2}{\beta_2} \dots \right) = \frac{x^a y^b}{B(a,b)} \left(\frac{1}{\beta_1} \frac{\alpha_2}{\beta_2} \dots \right),$$

which is relation (15). The expressions (15.4) and (15.5) for computing (15) follow from Theorem 2 [1; p.19], where $\lim_{n \rightarrow \infty} \frac{A_n}{B_n} = \frac{\alpha_1}{\beta_1} \frac{\alpha_2}{\beta_2} \dots$.

If $n \leq b$, then $\alpha_n \geq 0$ and β_n is a positive value not near 0. To insure that the maximum number of iterations satisfies $n \leq b$, (15) is applied only when $b \geq 40$. In this case, x must also be a sufficient distance from p when $a > 100$.

BASYM

Let γ be an arbitrary positive scaling factor that is assigned below. Then

$$I_x(a,b) \approx U \frac{2}{\sqrt{\pi}} e^{-z^2} \sum_{n=0}^N d_n J_n(z) (\beta \gamma)^n, \quad x \leq p \quad (16)$$

$$p = \frac{a}{a+b}, \quad q = \frac{b}{a+b}, \quad \beta = \sqrt{\frac{ab}{(a+b)^3}}, \quad U = \frac{p^a q^b}{B(a,b)} \sqrt{\frac{2\pi(a+b)}{ab}}$$

$$\phi(t) = t - 1 - \ln t, \quad t > 0 \quad (16.1)$$

$$\varphi(t) = a\phi\left(\frac{t}{p}\right) + b\phi\left(\frac{1-t}{q}\right), \quad 0 < t < 1$$

$$z = \sqrt{\varphi(x)}$$

$$a_n = \gamma^{-n} \lambda_n, \quad \gamma > 0 \quad (16.2)$$

$$\lambda_n = \frac{2}{n+2} [q p^{-n} + (-1)^n p q^{-n}], \quad (\lambda_0 = 1)$$

$$b_0^{(r)} = 1 \quad (16.3)$$

$$b_n^{(r)} = r a_n + \frac{1}{n} \sum_{i=1}^{n-1} [r(n-i)-i] b_i^{(r)} a_{n-i}, \quad n=1,2,\dots \quad [11]$$

$$c_n = \frac{1}{n} b_{n-1}^{(-n/2)}, \quad (n \geq 1)$$

$$d_0 = 1, \quad d_n = - \sum_{i=0}^{n-1} d_i c_{n-i+1} \quad (16.4)$$

$$J_n(z) = 2^{\frac{n}{2}-1} e^{-z^2} \int_z^\infty e^{-v^2} v^n dv. \quad (16.5)$$

$$\beta \gamma = \sqrt{q/a}, \quad a \leq b \quad (16.6)$$

$$\beta \gamma = \sqrt{p/b}, \quad a \geq b.$$

N is defined as the smallest integer such that

$$|d_N J_N(z) (\beta \gamma)^N| + |d_{N-1} J_{N-1}(z) (\beta \gamma)^{N-1}| \leq c \sum_{n=0}^N d_n J_n(z) (\beta \gamma)^n.$$

Relation (16) is used only when a and b are large and $x \approx p$. It appears to be new. Given $\lambda = (a+b)(p-x)$ and a tolerance ϵ , (16) is computed by the function BASYM($a, b, x, y, \lambda, \epsilon$). The expansion is obtained by writing (1) in the form

$$I_x(a, b) = \frac{p^a q^b}{B(a, b)} \int_0^x \left(\frac{t}{p}\right)^a \left(\frac{1-t}{q}\right)^b \frac{dt}{t(1-t)}, \quad 0 < x < 1. \quad (16.7)$$

From (16.1), $\varphi(t) > 0$ for $t \neq p$ and

$$\varphi(x) = -\left(a \ln \frac{x}{p} + b \ln \frac{y}{q}\right) \quad (16.8)$$

Thus if $u = \sqrt{\varphi(t)}$ for $t \leq p$ and $u = -\sqrt{\varphi(t)}$ for $t > p$, then

$$I_x(a, b) = U \beta \sqrt{\frac{2}{\pi}} \int_z^\infty e^{-u^2} \frac{u}{p-t} du, \quad (16.9)$$

where $z = \sqrt{\varphi(x)}$ if $x \leq p$ and $z = -\sqrt{\varphi(x)}$ if $x > p$. Hereafter, we shall assume that $x \leq p$. In order to use (16.9), $u/(p-t)$ will be replaced by its Maclaurin series in u .

From (16.8) and

$$\begin{aligned} a \ln \frac{t}{p} &= a \ln \left(1 - \frac{p-t}{p}\right) = -a \sum_{n=1}^{\infty} \frac{1}{n} \left(\frac{p-t}{p}\right)^n \\ b \ln \frac{1-t}{q} &= b \ln \left(1 + \frac{p-t}{q}\right) = -b \sum_{n=1}^{\infty} \frac{1}{n} \left(\frac{t-p}{q}\right)^n, \end{aligned}$$

one obtains

$$u^2 = \varphi(t) = \frac{1}{2\beta^2} (p-t)^2 \sum_{n \geq 0} \lambda_n (p-t)^n, \quad |p-t| < \min\{p, q\},$$

where λ_n is given in (16.2). Thus if $a_n = \gamma^{-n} \lambda_n$ for $\gamma > 0$, then $2(\beta \gamma u)^2 = s^2 A$, where $A = \sum_{n \geq 0} a_n s^n$ and $s = \gamma(p-t)$. Hence, if $P(s) = s\sqrt{A}$ then $\sqrt{2} \beta \gamma u = P(s)$. If $v = P(s)$, since P is analytic at 0 with $P(0) = 0$, by the Lagrange-Bürmann expansion [8; p.58] the inverse $s = P^{-1}(v)$ of P is given by

$$\begin{aligned} s &= \sum_{n \geq 1} c_n v^n, \\ c_n &= \frac{1}{n} \text{res}(P^{-n}) = \frac{1}{n} \text{res}(s^{-n} A^{-n/2}), \end{aligned}$$

where $\text{res}(P^{-n})$ is the residue of the series P^{-n} . Also for any $r \neq 0$

$$A^r = \sum_{k \geq 0} b_k^{(r)} s^k,$$

where $b_k^{(r)}$ is given in (16.3), [11]. Consequently,

$$c_n = \frac{1}{n} b_{n-1}^{(-n/2)}$$

$$\sqrt{2} \beta \frac{u}{p-t} = \frac{v}{s} = 1 / \sum_{n \geq 1} c_n v^{n-1} = \sum_{n \geq 0} d_n v^n,$$

where d_n is given in (16.4), [11].

Now if $h=a/b$ for $a \leq b$ and $h=b/a$ for $a > b$, let $\gamma=(1+h)/h$, (see (16.6)). Then $A = \sum_{n \geq 0} a_n s^n$ for $|s| < 1$ and

$$\begin{aligned} a_n &= \frac{2}{n+2} q \left[1 + (-1)^n h^{n+1} \right], \quad a \leq b \\ a_n &= \frac{2}{n+2} p \left[(-1)^n + h^{n+1} \right], \quad a > b. \end{aligned} \quad (16.10)$$

It can be shown for $a=b$ that the series $v/s = \sum_{n \geq 0} d_n v^n$ has a radius of convergence no larger than $\sqrt{2\pi}$.

Indeed, we have

$$u^2 = av^2 = -a \ln(1-s^2),$$

and, with v complex, $s=0$ when $v=\sqrt{2\pi} \exp(i\pi/4)$.

If $a \leq b$ then $A = \sum_{n \geq 0} a_n s^n \geq 0$ for $0 \leq s < 1$ since each $a_n \geq 0$. Hence $v=s\sqrt{A}$ is defined and real for $0 \leq s < 1$. Also, from (16.8) we note that $v \rightarrow \infty$ when $s \rightarrow 1$, and $dv/ds = (s/v)[(1-s)(1+hs)]^{-1} \neq 0$ for $0 < s < 1$. Consequently, $f(v) = v/s = \sqrt{A}$ can be regarded as a function of v for all $v \geq 0$. Since it is also true that the derivatives $f^{(n)}(v)$ are bounded for $v \geq 0$ (see Appendix C), let $M_n = \sup\{|f^{(n)}(v)| : v \geq 0\}$ for $n \geq 1$. Then $M_1 \leq 1$ (see Appendix C), and for

$$\psi_n(v) = f(v) - \sum_{i=0}^{n-1} d_i v^i,$$

$|\psi_n(v)| \leq M_n v^n / n!$ ($v > 0$) by the Taylor formula with remainder. Therefore, from $\sqrt{2}\beta u/(p-t) = f(v)$ and (16.9) we obtain

$$I_X(a,b) = U \frac{2}{\sqrt{\pi}} e^{-z^2} \left[\sum_{i=0}^{n-1} d_i (\beta\gamma)^i J_i(z) + E_n \right], \quad z \geq 0$$

$$E_n = \frac{1}{2} e^{z^2} \int_z^\infty e^{-u^2} \psi_n(\sqrt{2}\beta\gamma u) du,$$

where $J_n(z)$ is given by (16.5). Also

$$|E_n| \leq M_n / n! (\beta\gamma)^n J_n(z) = M_n / n! J_n(z) [a(1+h)]^{-n/2} \quad (16.11)$$

so that $|E_n| \rightarrow 0$ when $a \rightarrow \infty$ for fixed b . Consequently, (16) is asymptotic for $a \leq b$. Also we note that

$$J_n(z) = 2^{(n/2)-2} e^{z^2} \Gamma[(n+1)/2] Q[(n+1)/2, z^2],$$

where Q is defined by (14.3), and that $J_n(z)$ can be computed recursively by

$$J_0(z) = (\sqrt{\pi}/4) e^{z^2} \operatorname{erfc}(z), \quad J_1 = \alpha^{-3/2}$$

$$J_n(z) = 2^{-3/2} (\sqrt{2} z)^{n-1} + (n-1) J_{n-2}(z), \quad n \geq 2.$$

When $a > b$ the situation is less satisfactory, since it has not yet been shown that (16) is asymptotic. Nevertheless, the utility of (16) has been established for $y < 1.05q$ when $b \geq 100$ by extensive computer testing.

Finally, we observe that the definition of U given in (16) is not suitable for computational purposes. U can be accurately evaluated using

$$\Delta(a) = \ln \Gamma(a) - (a - \frac{1}{2}) \ln a + a - \frac{1}{2} \ln(2\pi) \quad (16.12)$$

$$\ln U = \Delta(a+b) - \Delta(a) - \Delta(b).$$

III. DOMAINS FOR SUBPROGRAMS

In this section we specify the regions of application for the five subprograms discussed in the previous section. A flowchart is given in Fig. 1.

In order to establish such regions the following conditions must be met:

- (a) The applied algorithms must be efficient and yield the desired accuracy over such regions.
- (b) It is necessary that $J \leq .9$, where J has the value $I_X(a,b)$ or $I_Y(b,a)$. J is computed by the proper choice, according to (a), of one or more of the subprograms based on (12)–(16). The complement is then obtained as $1-J$.

Even though analysis was carried out to predict the efficiency and achievable accuracy of the basic relations over various domains, (a) was established with exhaustive testing by Morris using double precision versions of the subprograms.

Since J is to be computed to the greatest possible accuracy, the relative tolerance to be satisfied will be $\epsilon = \max\{\epsilon_0, 10^{-15}\}$ where ϵ_0 is the smallest number for which $1 + \epsilon_0 > 1$ for the floating point arithmetic being used. The restriction that $\epsilon \geq 10^{-15}$, thereby limiting the maximum precision to 14–15 significant digits, is made since many of the supporting subprograms are accurate to a maximum of 14 digits (see section IV).

Arguments are presented near the end of this section showing that condition (b) is always satisfied. For easy reference we have:

BPSEN(a,b,x, ϵ)	Relation(12)
BUP(a,b,x,y,N, ϵ)	Relation(13)
BGRAT(a,b,x,y,w,15 ϵ ,IERR)	Relation(14)
BFRAC(a,b,x,y, λ ,15 ϵ)	Relation(15)
BASYM(a,b,x,y, λ ,100 ϵ)	Relation(16)

There are two main domains to consider, namely $\min(a,b) \leq 1$ and $\min(a,b) > 1$. It should be recalled that if a and b , and x and y are interchanged in one of the above subprograms, then the subprogram, except for BUP, yields $J = I_Y(b,a) = 1 - I_X(a,b)$ with $1 - J = I_X(a,b)$. BUP gives on the interchange $I_Y(b,a) - I_Y(b+N,a)$.

$\min(a,b) \leq 1$

If $x > 1/2$, then a and b , and x and y are interchanged. For $x \leq 1/2$, (17) is used for computing $J = I_x(a, b)$ and (18)–(20) are used for computing $J = I_y(a, b)$. In (19) and (20), w_0 is the initial value of w and J is the final value of w .

$$\text{BPSER}(a, b, x, \epsilon) \quad \max(a, b) \leq 1, \quad a \geq \min(0.2, b) \quad (17)$$

$$\max(a, b) \leq 1, \quad a < \min(0.2, b), \quad x^a \leq 0.9 \quad (17.1)$$

$$\max(a, b) > 1, \quad b \leq 1 \quad (17.2)$$

$$\max(a, b) > 1, \quad b > 1, \quad x < 0.1, \quad (bx)^a \leq 0.7 \quad (17.3)$$

$$\text{BPSER}(b, a, y, \epsilon) \quad \max(a, b) \leq 1, \quad a < \min(0.2, b), \quad x^a > 0.9, \quad x \geq 0.3 \quad (18)$$

$$\max(a, b) > 1, \quad b > 1, \quad x \geq 0.3 \quad (18.1)$$

$$\text{BGRAT}(b, a, y, x, w, 15\epsilon, \text{IERR}), \quad w_0 = 0$$

$$\max(a, b) > 1, \quad b > 15, \quad 0.1 \leq x < 0.3 \quad (19)$$

$$\max(a, b) > 1, \quad b > 15, \quad x < 0.1, \quad (bx)^a > 0.7 \quad (19.1)$$

$$\text{BGRAT}(b + N, a, y, x, w, 15\epsilon, \text{IERR}), \quad w_0 = \text{BUP}(b, a, y, x, N, \epsilon), \quad N = 20$$

$$\max(a, b) > 1, \quad b > 1, \quad 0.1 \leq x < 0.3, \quad b \leq 15 \quad (20)$$

$$\max(a, b) > 1, \quad b > 1, \quad x < 0.1, \quad (bx)^a > 0.7, \quad b \leq 15 \quad (20.1)$$

$$\max(a, b) \leq 1, \quad a < \min(0.2, b), \quad x^a > 0.9, \quad x < 0.3 \quad (20.2)$$

$\min(a,b) > 1$

If $x > p$ then a and b , and x and y are interchanged. For $x \leq p$, (21)–(26) are used for computing $J = I_x(a, b)$. In (22)–(24) N is the largest integer less than b and $\bar{b} = b - N$. Also, in (23) and (24) w_0 is the initial value of w and J is the final value of w .

$$\text{BPSER}(a, b, x, \epsilon) \quad b < 40, \quad bx \leq .7 \quad (21)$$

$$\text{BUP}(\bar{b}, a, y, x, N, \epsilon) + \text{BPSER}(a, \bar{b}, x, \epsilon) \quad b < 40, \quad bx > .7, \quad x \leq .7 \quad (22)$$

$$\text{BGRAT}(a, \bar{b}, x, y, w, 15\epsilon, \text{IERR}), \quad w_0 = \text{BUP}(\bar{b}, a, y, x, N, \epsilon) \quad b < 40, \quad x > .7, \quad a > 15 \quad (23)$$

$$\begin{aligned}
& \text{BGRAT}(a+M, \bar{b}, x, y, w, 15\epsilon, \text{IERR}), \quad M=20 \\
& w_0 = \text{BUP}(\bar{b}, a, y, x, N, \epsilon) + \text{BUP}(a, \bar{b}, x, y, M, \epsilon) \\
& \quad b < 40, \quad x > .7, \quad a \leq 15
\end{aligned} \tag{24}$$

$$\begin{aligned}
& \text{BASYM}(a, b, x, y, \lambda, 100\epsilon) \quad b \geq 40, \quad 100 < a \leq b, \quad x \geq .97p \\
& \quad b \geq 40, \quad 100 < b < a, \quad y \leq 1.03q
\end{aligned} \tag{25}$$

(25.1)

$$\begin{aligned}
& \text{BFRAC}(a, b, x, y, \lambda, 15\epsilon) \quad b \geq 40, \quad a \leq b, \quad a \leq 100
\end{aligned} \tag{26}$$

(26.1)

$$\begin{aligned}
& b \geq 40, \quad 100 < a \leq b, \quad x < .97p \\
& b \geq 40, \quad a > b, \quad b \leq 100
\end{aligned} \tag{26.2}$$

(26.3)

$$\begin{aligned}
& b \geq 40, \quad 100 < b < a, \quad y > 1.03q
\end{aligned} \tag{26.3}$$

These statements are summarized in the flowchart for BRATIO in Fig.1. Proofs are now given which verify that $J \leq 0.9$ is always satisfied (requirement b above). The arguments use the facts that $\ln \Gamma(t)$ is strictly convex for $t > 0$, [1; 6.4.10], and that $I_X(a, b)$ is a decreasing function of a and an increasing function of b . The latter result is proven in Appendix B. Since $\ln \Gamma(t)$ is strictly convex for $t > 0$, we also note that $\psi(t) = \frac{d}{dt}\{\ln \Gamma(t)\}$, [1; 6.3.1], is an increasing function of t .

For (17): $a \leq 1, b \leq 1, x \leq 1/2, a \geq \min(.2, b)$

If $a \geq 0.2$ then

$$J = I_X(a, b) \leq I_X(a, 1) = x^a \leq (1/2)^2 = 0.8706.$$

If $a \geq b$ then

$$J = I_X(a, b) \leq I_X(a, a) \leq I_{1/2}(a, a).$$

Also $I_{1/2}(a, a) = 0.5$ from (4).

For (17.1): $a \leq 1, b \leq 1, x \leq 1/2, a < \min(.2, b), x^a \leq 0.9$

$$J = I_X(a, b) \leq I_X(a, 1) = x^a \leq 0.9$$

For (17.2): $a > 1, b \leq 1, x \leq 1/2$

$$J = I_X(a, b) \leq I_X(1, 1) = x \leq 1/2$$

For (17.3): $a \leq 1, b > 1, x < 0.1, (bx)^a \leq 0.7$

Integrating (1) by parts gives

$$\begin{aligned}
I_X(a, b) &= G(a, b) \left\{ \frac{x^a}{a} (1-x)^{b-1} + \frac{b-1}{a} \int_0^x t^a (1-t)^{b-2} dt \right\} \\
&\leq G(a, b) \left\{ \frac{x^a}{a} (1-x)^{b-1} + \frac{b-1}{a} x^a \int_0^x (1-t)^{b-2} dt \right\} = G(a, b) \frac{x^a}{a}.
\end{aligned}$$

Let

$$F(a,b) = \Gamma(a+b)/[\Gamma(b)b^a], \quad 0 \leq a \leq 1, \quad b \geq 1.$$

Since $\partial/\partial a [\ln F] = \psi(a+b) - \ln b$ is an increasing function of a for $a > 0$, $\ln F$ is strictly convex and hence F is strictly convex for $a > 0$. Thus $F(a,b) \leq 1$ for $0 \leq a \leq 1$ since $F(0,b) = F(1,b) = 1$. Hence

$$J = I_X(a,b) \leq (bx)^a / \Gamma(a+1) \leq 0.7 / \Gamma(1.46163\ldots) < 0.791.$$

For (18): $a \leq 1, b \leq 1, 0.3 \leq x \leq 0.5, a < \min(0.2, b), x^a > 0.9$

From (1), with

$$(1-t)^{b-1} \geq 1, \quad 0 \leq t \leq x,$$

$$I_X(a,b) \geq H(a) \frac{b}{a+b} x^a \geq H(a)(0.9)/2 \geq .45, \quad (a < b),$$

where

$$H(a) = \Gamma(a+b+1)/[\Gamma(a+1)\Gamma(b+1)] \quad (0 \leq a \leq 1).$$

The last inequality on I_X follows from the fact that $H(a)$ is increasing and that $H(0) = 1$. Hence

$$J = I_Y(b,a) = 1 - I_X(a,b) \leq 0.55.$$

For (18.1): $a \leq 1, b > 1, 0.3 \leq x \leq 0.5$

$$J = I_Y(b,a) < I_Y(1,1) = 1 - x \leq 0.7$$

For (19): $a \leq 1, b > 15, 0.1 \leq x < 0.3$

$$J = I_Y(b,a) \leq I_Y(b,1) = y^b = (1-x)^b \leq (0.9)^{15} \approx 0.2059$$

For (19.1): $a \leq 1, b > 15, x < 0.1, (bx)^a > 0.7$

Then

$$J = I_Y(b,a) \leq I_Y(b,1) = y^b = (1-x)^b.$$

From (1), using $(1-t)^{b-1} \geq (1-x)^{b-1}$ and recalling $F(a,b)$ in the proof of (17.3),

$$I_X(a,b) \geq \frac{F(a,b)}{\Gamma(a)} \frac{(1-x)^b}{1-x} \frac{(bx)^a}{a} \geq F(a,b)(bx)^a J / \Gamma(a+1).$$

Now $\partial F / \partial b = g(a,b)F$ where $g(a,b) = \psi(a+b) - \psi(b) - a/b$. Since $g_{aa} < 0$ [1; 6.4.10], g is strictly concave for $0 < a < 1$. Thus, since $g(0,b) = g(1,b) = 0$ [1; 6.3.5], $g(a,b) > 0$ for $0 < a < 1$ and $\partial F / \partial b > 0$. Hence, F is increasing in b for $0 < a < 1$ and

$$F(a,b) > F(a,1) = \Gamma(a+1).$$

Thus $I_X(a,b) \geq 0.7J$ for $0 \leq a \leq 1$, which implies that $J \leq 1/1.7 \approx .59$.

For (20): $a \leq 1, 1 < b \leq 15, 0.1 \leq x < 0.3$

$$J = I_y(b,a) \leq I_y(b,1) = y^b = (1-x)^b \leq (0.9)^b \leq 0.9$$

For (20.1): $a \leq 1, 1 < b \leq 15, x < 0.1, (bx)^a > 0.7$

Proof is the same as that given for (19.1).

For (20.2): $b \leq 1, x < 0.3, a < \min(2, b), x^a > 0.9$

$$I_x(a,b) \geq I_x(a,a) = K(a) \frac{a^2}{2a} \int_0^x t^{a-1} (1-t)^{a-1} dt,$$

where

$$K(a) = \Gamma(2a+1)/[\Gamma(a+1)\Gamma(a+1)], \quad (0 \leq a \leq 1).$$

From (i), using

$$(1-t)^{a-1} \geq 1, \quad 0 \leq t \leq x,$$

it follows that

$$I_x(a,b) > K(a) x^a / 2 > 0.45 K(a).$$

Since $\partial K(a)/\partial a = 2K(a)[\psi(2a+1) - \psi(a+1)] > 0$, $K(a)$ is an increasing function. Hence $K(a) \geq 1$ and

$$J = I_y(b,a) \leq 0.55.$$

For (21)–(26): $a > 1, b > 1$

In this case it can be shown that if $x \leq p$ then

$$I_x(a,b) \leq \begin{cases} 1 - 1/e & a < b \\ 1/2 & a \geq b. \end{cases} \quad (27)$$

A proof of this result is given in Appendix A. Hence

$$J = \begin{cases} I_x(a,b) & x \leq p \\ I_y(b,a) & x > p. \end{cases} \quad (28)$$

Notation for the flowchart in Fig.1

- 12 Refers to (12) for computing $I_x(a,b)$ (BPSER)
12̄ Refers to (12) for computing $I_y(b,a)$ (BPSER)
13 Refers to (13) for computing $I_x(a,b)$ (BUP)
13̄ Refers to (13) for computing $I_y(b,a)$ (BUP)
14 Refers to (14) for computing $I_x(a,b)$ (BGRAT)
14̄ Refers to (14) for computing $I_y(b,a)$ (BGRAT)
15 Refers to (15) for computing $I_x(a,b)$ (BFRAC)
15̄ Refers to (15) for computing $I_y(b,a)$ (BFRAC)
16 Refers to (16) for computing $I_x(a,b)$ (BASYM)
16̄ Refers to (16) for computing $I_y(b,a)$ (BASYM)

$$p = a/(a+b), \quad q = b/(a+b)$$

I.C. \Rightarrow Interchange a and b; x and y.

$[b] \equiv$ largest integer $< b$.

$$\lambda \equiv a - (a+b)x, \quad a \leq b; \quad \lambda \equiv (a+b)y - b, \quad a > b.$$

Numbers above some of the flowchart boxes refer to the labels in the Fortran listing of BRATIO.




Fig. 1 Flowchart for BRATIO (Notation on p. 16)

IV. AUXILIARY FUNCTIONS

In order to compute $I_x(a,b)$ and $1 - I_x(a,b)$, procedures are needed for evaluating $\Gamma(a)$, $\ln \Gamma(a)$, the error function $\operatorname{erf} x$, $\exp(x^2) \operatorname{erfc} x$, the incomplete gamma function $Q(a,x)$, (14.3), for $a \leq 1$, and the functions

$$\begin{aligned}
 & e^x - 1 && (29) \\
 & \ln(1+x) && (|x| \leq .375) \\
 & \ln \Gamma(1+x) && (-0.2 \leq a \leq 1.25) \\
 & 1/\Gamma(1+a) - 1 && (-0.5 \leq a \leq 1.5) \\
 & e^{-x} x^a / \Gamma(a) && (a > 0, x \geq 0) \\
 & \phi(x) = x - 1 - \ln x && (x > 0).
 \end{aligned}$$

These functions are discussed in [6]. Also, procedures are needed for computing

$$\begin{aligned}
 \Delta(a) &= \ln \Gamma(a) - (a - .5) \ln a + a - .5 \ln(2\pi) && (a \geq 8) \\
 \text{ALGDIV}(a,b) &= \ln [\Gamma(b)/\Gamma(a+b)], && (a \geq 0, b \geq 8) \\
 \text{BCORR}(a,b) &= \Delta(a) + \Delta(b) - \Delta(a+b) && (a, b \geq 8) \\
 \text{BETALN}(a,b) &= \ln B(a,b) && (a, b > 0) \\
 \text{BRCOMP}(a,b,x,y) &= x^a y^b / B(a,b) && (a, b > 0, 0 < x < 1, y = 1-x).
 \end{aligned} \tag{30}$$

Rational minimax approximations are used for the functions given in (29). Experience indicates that such approximations normally generate less error and can be considerably more efficient than the standard expansions. However, minimax approximations have the disadvantage of being limited to a fixed maximum precision. The minimax approximations used are designed to achieve a maximum precision of 14 significant digits.

If $\Delta(a)$ is needed only for $a \geq 20$, then the sum

$$1/(12a) - 1/(360a^3) + \dots$$

in the asymptotic expansion of $\ln \Gamma(a)$ [1; 6.1.41] may be used. If $a \geq 15$ then the minimax approximation

$$\Delta(a) = \sum_{n=0}^4 c_n / a^{2n+1}$$

$$c_0 = .83333\ 33333\ 33333E-01$$

$$c_1 = -.27777\ 77777\ 70481E-02$$

$$c_2 = .79365\ 06631\ 83693E-03$$

$$c_3 = -.59515\ 63364\ 28591E-03$$

$$c_4 = .82075\ 63703\ 53826E-03$$

can be applied, and if $a \geq 8$ the minimax approximation

$$\Delta(a) = \sum_{n=0}^5 d_n / a^{2n+1}$$

$$d_0 = .83333\ 33333\ 33333E-01 \quad (32)$$

$$d_1 = -.27777\ 77777\ 60991E-02$$

$$d_2 = .79365\ 06668\ 25390E-03$$

$$d_3 = -.59520\ 29313\ 51870E-03$$

$$d_4 = .83730\ 80340\ 31215E-03$$

$$d_5 = -.16532\ 29627\ 80713E-02$$

can be used. These approximations were obtained by Morris [9]. On the CDC 6000-7000 series computers, they are accurate to within 1 unit of the 14th significant digit.

Expansions for ALGDIV(a,b) and BCORR(a,b) use (32). From the definition of Δ

$$\text{ALGDIV}(a,b) = w - (a+b-.5)\ln(1+a/b) - a(\ln b - 1) \quad (33)$$

$$w = \Delta(b) - \Delta(a+b).$$

Let

$$p = a/(a+b), \quad q = b/(a+b), \quad S_m = 1 + q + \dots + q^{m-1} \quad (m \geq 1).$$

Then

$$1 - q^m = (1 - q)S_m = pS_m, \quad \frac{pS_m}{b^m} = \frac{1}{b^m} - \frac{1}{(a+b)^m}. \quad (34)$$

Thus, from (32) we obtain

$$w = \frac{p}{b} \sum_{n=0}^5 d_n \frac{S_{2n+1}}{b^{2n}}, \quad (35)$$

which completes the algorithm for ALGDIV(a,b). Also

$$\text{BCORR}(a,b) = \Delta(a_0) + [\Delta(b_0) - \Delta(a_0 + b_0)]$$

$$a_0 = \min\{a,b\}, \quad b_0 = \max\{a,b\},$$

where (32) and (35) are applied.

If $a \leq b$, then BETALN(a,b) can be accurately computed when $a \geq 1$. If $a \geq 8$ then

$$\text{BETALN}(a,b) = (.5 \ln(2\pi) - .5 \ln b) + \text{BCORR}(a,b) - u - v$$

$$u = -(a - .5) \ln[a/(a+b)], \quad v = b \ln(1+a/b)$$

is applied. If $2 < a < 8$ then a is reduced to the interval $[1,2]$ by

$$B(a,b) = \frac{a-1}{a+b-1} B(a-1,b).$$

Consequently, it can be assumed that $a \leq 2$.

If $b \geq 8$ then

$$\ln B(a,b) = \ln \Gamma(a) + ALGDIV(a,b)$$

is applied. If $2 < b < 8$ then b is also reduced to the interval $[1,2]$ when $a \geq 1$. Thus, we need only consider the cases: $1 \leq a \leq 2$, $1 \leq b \leq 2$, or $a < 1$ and $b < 8$. If $a \geq 1$ then

$$\ln B(a,b) = \ln \Gamma(a) + \ln \Gamma(b) - \ln \Gamma(a+b)$$

is appropriate. No loss of accuracy due to subtraction can occur since $\ln \Gamma(a)$, $\ln \Gamma(b)$, and $-\ln \Gamma(a+b)$ are nonpositive. However, subtraction does occur when $a < 1$. It currently is not clear how loss of accuracy due to subtraction can be avoided when $a < 1$. Therefore, in this case BETALN is not used in BRATIO.

If $\min\{a,b\} < 8$ then BRCOMP(a,b,x,y) can be computed directly from its definition. Otherwise,

$$\begin{aligned} BRCOMP(a,b,x,y) &= \sqrt{\frac{ab}{2\pi(a+b)}} e^{-z} \\ z &= [a\phi(1 - \lambda/a) + b\phi(1 + \lambda/b)] + BCORR(a,b) \end{aligned}$$

is used, where λ is given in (15.1).

V. CONCLUDING REMARKS

Formulas (12), (13), (14), (15), and (16) for $I_x(a,b)$ are of the form γS , where S is a series. For example, for (15) $\gamma = x^a y^b / B(a,b)$ and S is a continued fraction. On the CDC 6000-7000 series computers, almost no error is generated in computing the series S over the domains specified in section III. The series is normally accurate to within 1 or 2 units of the 14th significant digit. However, the precision of the factor γ is restricted by the inherent error of $I_x(a,b)$. Extensive testing on the CDC 6000-7000 series computers comparing the results obtained by BRATIO with results from double precision code, indicates that the precisions of the values obtained for $I_x(a,b)$ and $1 - I_x(a,b)$ by BRATIO approximate the inherent errors of these functions up to a maximum of 14 significant digits. On any computer, accuracy is restricted to 14 digits because of the algorithms used for the auxiliary functions in section IV.

On the CDC 6000-7000 series computers, a maximum of 7 terms of the series (14) for BGRAT, and a maximum of 11 terms of (16) for BASYM were observed for the domains specified in section III. Frequently, 40 or fewer terms of (12) for BPSER suffice, but a maximum of 92 terms has been observed when a is small, b is large, and $x \approx .3$. Also, 40 or fewer terms generally suffice for the continued fraction (15), BFRAC, but a maximum of 58 terms has been observed when a or b is exceedingly large and $x \approx a/(a+b)$.

In practice, BRATIO has been found to be a reliable and efficient subroutine. As was noted in the previous sections, in order to develop such a subroutine new formulas were needed for $I_x(a,b)$, a surprisingly elaborate specification of the domains of usage for the various formulas had to be given, and a number of auxiliary functions had to be treated with extreme care. Thus, the development of BRATIO for efficiently computing $I_x(a,b)$ and $1 - I_x(a,b)$ to high relative accuracy was not a simple task.

VI. NUMERICAL EXAMPLES

A collection of 16 examples using BRATIO is given in Fig.2. The results were obtained using the CDC 6000-7000 series single precision truncation floating arithmetic. As was noted in section III, the function BUP is used only with BPSER or BGRAT and appears in (20), (22), (23), and (24). The following three cases illustrate its use. The quantity ϵ below is set to approximately .710E-14 (see p.11).

Case (1). $a=.10, b=14.5, x=.29, y=.71$

$$\begin{aligned} \text{From (20)} \quad w_0 &= \text{BUP}(14.5, .10, .71, .29, 20, \epsilon) \\ &= I_{.71}(14.5, .10) - I_{.71}(34.5, .10) \\ &= .17776\ 09989\ 0838E-3. \end{aligned}$$

Hence, if w is assigned the initial value w_0 then a call to BGRAT(34.5, .10, .71, .29, w , 15 ϵ , IERR) yields the value

$$\begin{aligned} w &= w_0 + I_{.71}(34.5, .10) = I_{.71}(14.5, .10) \\ &= .17785\ 31648\ 7898E-3. \end{aligned}$$

Also

$$I_{.29}(.10, 14.5) = .99982\ 21468\ 3512.$$

Case (2). $a=1.5, b=20.5, x=.065, y=.935$

(Note that $bx > .70, \lambda = a - (a+b)x = .07 > 0$)

$$\begin{aligned} \text{From (22)} \quad w_0 &= \text{BUP}(.50, 1.5, .935, .065, 20, \epsilon) \\ &= I_{.935}(.50, 1.5) - I_{.935}(20.5, 1.5) \\ &= .56745\ 07805\ 9439. \end{aligned}$$

Then

$$\begin{aligned} I_{.065}(1.5, 20.5) &= w_0 + \text{BPSER}(1.5, .50, .065) \\ &= w_0 + .71754\ 32115\ 7741E-3 \\ &= .57462\ 62127\ 1016, \end{aligned}$$

$$I_{.935}(20.5, 1.5) = .42537\ 37872\ 8984.$$

Case (3). $a=10.5, b=1.5, x=.80, y=.20, (\lambda > 0)$.

$$\begin{aligned} \text{From (24)} \quad \bar{w}_0 &= \text{BUP}(.50, 10.5, .20, .80, 1, \epsilon) \\ &= I_{.20}(.50, 10.5) - I_{.20}(1.5, 10.5) \\ &= .15518\ 20005\ 6352, \\ w_0 &= \bar{w}_0 + \text{BUP}(10.5, .50, .80, .20, 20, \epsilon) \\ &= \bar{w}_0 + I_{.80}(10.5, .50) - I_{.80}(30.5, .50) \\ &= \bar{w}_0 + .32149\ 19363\ 8971E-1 \\ &= .18733\ 11942\ 0245. \end{aligned}$$

Hence, if w is assigned the initial value w_0 then a call to BGRAT(30.5, .50, .80, .20, w, 15 ϵ , IERR) yields the value

$$w = w_0 + I_{.80}(30.5, .50)$$

or

$$I_{.80}(10.5, 1.5) = .18756\ 94122\ 3880.$$

Also,

$$I_{.20}(1.5, 10.5) = .81243\ 05877\ 6120.$$

a	b	x	y	$I_x(a,b)$	$1 - I_x(a,b)$
.1	.8	.40	.60	.88776 70523 5302E+00	.11223 29476 4698E+00
.1	.8	.60	.40	.92957 83432 6833E+00	.70421 65673 1668E-01
.1	2.3	.40	.60	.97448 97683 7361E+00	.25510 23162 6386E-01
.1	2.3	.60	.40	.99196 58486 2884E+00	.80341 51371 1598E-02
5.0	40.0	.99	.01	.10000 00000 0000E+01	.13053 04681 1410E-74
5.0	10.0	.99	.01	.10000 00000 0000E+01	.96509 74271 4997E-17
10.0	38.0	.02	.98	.26944 43561 3309E-07	.99999 99730 5556E+00
70.0	10.0	.85	.15	.23472 44941 6827E+00	.76527 55058 3173E+00
* 70.0	50.0	.99	.01	.10000 00000 0000E+01	.54279 07073 1686E-66
* 70.0	50.0	.10	.90	.47438 77486 2163E-38	.10000 00000 0000E+01
* 75.0	50.0	.10	.90	.61550 21193 1591E-42	.10000 00000 0000E+01
* 500.0	501.0	.60	.40	.99999 99999 3299E+00	.67009 77013 4757E-10
500.0	501.0	.40	.60	.10148 03038 4399E-09	.99999 99998 9852E+00
1000.0	1001.0	.49	.51	.19153 11043 9543E+00	.80846 88956 0457E+00
1001.0	1000.0	.49	.51	.17957 42144 6754E+00	.82042 57855 3246E+00
$(a=5.0E+20, \ b=5.0E+3, \ x=1.0, \ y=1.0E-17)$ $I_x(a,b)=.49811\ 93659\ 6617, \ I_y(b,a)=.50188\ 06340\ 3383)$					

Due to inherent error, the 4 starred examples are correct to within 1 unit of the 12th significant digit. All other cases are correct to within 5 units of the 14th significant digit.

Fig. 2 16 Examples of BRATIO

REFERENCES

1. Abramowitz,M. and Stegun,I.(EDS.), *Handbook of Mathematical Functions*, Dover, N.Y., 1965.
2. Amos, D.E. and Daniel, S.L., Significant Digit Incomplete Beta Ratios Development, Report SC-DR-69-591, Sandia Laboratories, Albuquerque, N.M., 1969.
3. Aroian, L.A, Continued Fractions for the Incomplete Beta Function, *Am. Math. Statist.*, 12, 1941, 218-223.
4. DiDonato, A.R. and Jarnagin, M.P. Jr., A Method for Computing the Incomplete Beta Function Ratio, NWL Report No. 1949, U.S. Naval Weapons Laboratory, Dahlgren, Va. 22448, 1966.
5. DiDonato, A.R. and Jarnagin, M.P., The Efficient Calculation of the Incomplete Beta Function Ratio for Half - Integer Values of the Parameters a,b, *Math. of Comp.*, 21, 1967, 652-662.
6. DiDonato, A.R. and Morris, A.H., Computation of the Incomplete Gamma Function Ratios and Their Inverse, *ACM Trans. of Math. Software*, 12, 1986, 377-393.
7. Dutka, J., The Incomplete Beta Function - A Historical Profile, *Archive for History of Exact Sciences*, 24, 1981, 11-29.
8. Henrici P., *Applied and Computational Complex Analysis (Vol. 1)*, John Wiley and Sons, 1974.
9. Morris A.H., NSWC Library of Mathematical Subroutines, Rep. NSWC TR 86-251. Naval Surface Weapons Center, Dahlgren, Va. 22448, 1987.
10. Olver F.W.J., *Asymptotics and Special Functions*, Academic Press, 1974.
11. Pourahmadi, M.; Taylor Expansion of $\text{Exp}(\sum_{k=0}^{\infty} a_k z^k)$ and Some Applications, *American Math Monthly*, 1984, 303-307.
12. Temme, N.M., Uniform Asymptotic Expansions of the Incomplete Gamma Functions and the Incomplete Beta Function, *Math. of Comp.* 29, 1975, 1109-1114.
13. Tretter, M.J. and Walster, G.W., Continued Fractions for the Incomplete Beta Function: Additions and Corrections, *Ann. of Stat.*, 7, 1979, 462-465.
14. Wall, H.S., *Analytic Theory of Continued Fractions*. Van Nostrand Co., N.Y., 1948.
15. Wise, M.E., The Incomplete Beta Function as a Contour Integral and a Quickly Converging Series for Its Inverse, *Biometrika*, 37, 1950, 208-218.

APPENDIX A

BOUNDS ON $I_x(a,b)$ WHEN $x=a/(a+b)$, $\min\{a,b\} \geq 1$

BOUNDS ON $I_x(a,b)$ WHEN $x=a/(a+b)$, $\min\{a,b\} \geq 1$

The purpose of this appendix is to show that

$$\begin{aligned} 1/2 &\leq I_p(a,b) < 1 - e^{-1} && \text{if } 1 \leq a \leq b \\ 1/e &< I_p(a,b) \leq 1/2 && \text{if } 1 \leq b \leq a \end{aligned} \tag{A-1}$$

where $p=a/(a+b)$ and $q=1-p=b/(a+b)$. Since $I_{.5}(a,a)=.5$ from (4), we shall assume that $a \neq b$.

Lemma (A-1): If $\lambda(h)=(1+1/h)^{-h}$ for $h > 0$ then λ is a decreasing function. Also, $\lambda \rightarrow 1$ when $h \rightarrow 0$ and $\lambda \rightarrow 1/e$ when $h \rightarrow \infty$.

The lemma is given for reference. It is a well known result.

Corollary (A-1): If $a > 1$ then $1/e < I_p(a,1) < 1/2$.

Proof: From (1) we obtain $I_p(a,1)=p^a=(1+1/a)^{-a}$. Then from the lemma the corollary follows.

Corollary (A-2): If $b > 1$ then $1/2 < I_p(1,b) < 1 - 1/e$.

Proof: Immediate from (4) and corollary (A-1).

Since the above corollaries hold, in order to verify (A-1) it suffices to assume that $a, b > 1$. The following reasoning is due to James C. Perry (NSWC). Let

$$B \equiv \int_0^1 t^{a-1} (1-t)^{b-1} dt \tag{A-2}$$

$$B_p \equiv \int_0^p t^{a-1} (1-t)^{b-1} dt \tag{A-3}$$

$$\bar{B}_p \equiv \int_p^1 t^{a-1} (1-t)^{b-1} dt. \tag{A-4}$$

If $x=(1-t)^{b/a}$ then

$$B_p = \frac{a}{b} \int_{\lambda}^1 (x - x^{1+a/b})^{a-1} dx \tag{A-5}$$

$$\bar{B}_p = \frac{a}{b} \int_0^{\lambda} (x - x^{1+a/b})^{a-1} dx \tag{A-6}$$

where

$$\lambda = q^{b/a} = (1+a/b)^{-b/a}. \tag{A-7}$$

Also $1/e < \lambda < 1$ by lemma (A-1), and we now consider the functions

$$f(x) = x - x^{1+a/b} \text{ and } F(x) = f(x)^{a-1} \tag{A-8}$$

for $0 \leq x \leq 1$.

From (A-7) and (A-8) we note that

$$f(0) = F(0) = 0, \quad f(1) = F(1) = 0,$$

$f(x)$ and $F(x)$ have a unique maximum at $x = \lambda$, and

$f'(x)$ and $F'(x)$ are positive (negative) for $0 < x < \lambda$ ($\lambda < x < 1$).

For any x_2 such that $\lambda < x_2 \leq 1$ let

$$r(x_2) = \frac{x_2 - \lambda}{\lambda - x_1} \quad (\text{A-9})$$

where x_1 is the unique value in $[0, \lambda]$ where $f(x_1) = f(x_2)$ [see Fig. 3].

We now examine $r(x_2)$.




Fig. 3 Graph of $f(x)$

Lemma (A-2): If $g(x)$ is twice continuously differentiable on $[0, d]$, $g(0) = g'(0) = 0$, and g'' is increasing (decreasing) on $(0, d]$, then h is continuous and increasing (decreasing) on $[0, d]$ where

$$h(x) \equiv \begin{cases} g(x)/x^2 & (x > 0) \\ g''(0)/2 & (x = 0). \end{cases}$$

Proof: Assume that g'' is increasing on $(0, d]$. Since $h'(x) = (xg' - 2g)/x^3$ for $x > 0$, let $k(x) = xg' - 2g$ for $x \geq 0$. If $x > 0$ then there exists $0 < \xi < x$ such that $g'(x) = g''(\xi)x$ by the Mean-value theorem, so that

$$k'(x) = xg''(x) - g'(x) = x[g''(x) - g''(\xi)] > 0.$$

Therefore, $k(x)$ is increasing for $x > 0$. Since $k(0) = 0$, $h'(x) = k(x)/x^3 > 0$, which proves that h is increasing on $(0, d]$. Also, since $g(0) = g'(0) = 0$, $h(x) \rightarrow g''(0)/2$ when $x \rightarrow 0$ by L'Hopital's rule. If $g''(x)$ is decreasing on $(0, d]$ then apply the lemma to $-g$.

Theorem (A-1): Let $r(x)$ denote the function defined by (A-9). If $a < b$ ($a > b$) then $r(x)$ is increasing (decreasing) for $\lambda < x \leq 1$. Also, $r \rightarrow 1$ when $x \rightarrow \lambda$.

Proof: If $\phi(h) = f(\lambda) - f(\lambda + h)$ for $0 \leq h \leq 1 - \lambda$ then $\phi(0) = 0$, $\phi'(0) = -f'(\lambda) = 0$, and $\phi''(h) = (a/bq)(\lambda + h)^{-1+a/b}$ for $h \geq 0$. Hence, if $a < b$ ($a > b$) then ϕ'' is decreasing (increasing), so that

$\bar{\phi}(h) = \phi(h)/h^2$ is decreasing (increasing) by lemma (A-2). Also, $\bar{\phi}(h) \rightarrow \phi''(0)/2 = a/(2b\lambda)$ when $h \rightarrow 0$ by the lemma and $\lambda = q^{b/a}$.

Similarly, if $\psi(h) = f(\lambda) - f(\lambda - h)$ for $0 \leq h \leq \lambda$ then $\bar{\psi}(h) = \psi(h)/h^2$ is increasing (decreasing) when $a < b$ ($a > b$), and $\bar{\psi}(h) \rightarrow \psi''(0)/2 = a/(2b\lambda)$ when $h \rightarrow 0$.

Now if $\lambda < x \leq 1$ and $x_1 < \lambda$ where $f(x_1) = f(x)$, then

$$r^2 = \left(\frac{x-\lambda}{\lambda-x_1}\right)^2 = \frac{f(\lambda)-f(x_1)}{(\lambda-x_1)^2} \frac{(x-\lambda)^2}{f(\lambda)-f(x)} = \frac{\bar{\psi}(\lambda-x_1)}{\bar{\phi}(x-\lambda)}$$

so that $r^2 \rightarrow \frac{a/(2b\lambda)}{a/(2b\lambda)} = 1$ when $x \rightarrow \lambda$. If $a < b$ then when x increases, $\lambda - x_1$ increases, $\bar{\psi}(\lambda - x_1)$ increases, $\bar{\phi}(x - \lambda)$ decreases, and hence r^2 increases. Similarly, r^2 decreases when $a > b$. Consequently, since $r > 0$ the theorem follows.

Theorem (A-2): If $1 \leq a < b$ then $I_p(a,b) < 1 - 1/e$.

Proof: Since $a < b$, $r(x) \leq r(1) = (1-\lambda)/\lambda$ for $\lambda < x \leq 1$ by theorem (A-1). Also, since $I_p(1,b) < 1 - 1/e$ by corollary (A-2), we shall assume that $a > 1$. Now let

$$t = \frac{\lambda}{1-\lambda} (1-x) \quad \lambda \leq x \leq 1 \quad (\text{A-10})$$

so that $0 \leq t \leq \lambda$. For $x > \lambda$ let $x_1 < \lambda$ where $f(x_1) = f(x)$. Then $t < \lambda$, and from

$$r(x) = \frac{x-\lambda}{\lambda-x_1} \leq \frac{1-\lambda}{\lambda} \text{ we have } x_1 \leq t. \text{ Therefore,}$$

$$F\left(1 - \frac{1-\lambda}{\lambda} t\right) = F(x) = F(x_1) \leq F(t)$$

so that

$$B_p = \frac{a}{b} \int_{\lambda}^1 F(x) dx = \frac{a}{b} \frac{1-\lambda}{\lambda} \int_0^{\lambda} F\left(1 - \frac{1-\lambda}{\lambda} t\right) dt$$

$$\leq \frac{a}{b} \frac{1-\lambda}{\lambda} \int_0^{\lambda} F(t) dt = \left(\frac{1}{\lambda} - 1\right) \bar{B}_p < (e-1) \bar{B}_p$$

from (A-5) and (A-6). Therefore, from (A-2)–(A-4), $B_p < (e-1) \bar{B}_p = (e-1)(B - B_p)$ so that $I_p(a,b) = B_p/B < 1 - 1/e$.

Theorem (A-3): If $1 \leq b < a$ then $I_p(a,b) < 1/2$.

Proof: Since $a > b$, $r(x) < 1$ for $\lambda < x \leq 1$ by theorem (A-1). Also, since $I_p(a,1) < 1/2$ by corollary (A-1), we shall assume that $b > 1$. Now let

$$t = 2\lambda - x \quad \lambda \leq x \leq 1. \quad (\text{A-11})$$

Since $b/a < 1$, $.5 < \lambda < 1$ from (A-7) and lemma (A-1). Therefore, $0 < 2\lambda - 1 < \lambda$ and $2\lambda - 1 \leq t \leq \lambda$.

For $x > \lambda$ let $x_1 < \lambda$ where $f(x_1) = f(x)$. Then $t < \lambda$, and from $r(x) = \frac{x-\lambda}{\lambda-x_1} < 1$ we have $x_1 < t$.
 Therefore,

$$F(2\lambda-t) = F(x) = F(x_1) < F(t)$$

so that

$$\begin{aligned} B_p &= \frac{a}{b} \int_{\lambda}^1 F(x) dx = \frac{a}{b} \int_{2\lambda-1}^{\lambda} F(2\lambda-t) dt \\ &< \frac{a}{b} \int_{2\lambda-1}^{\lambda} F(t) dt \leq \frac{a}{b} \int_0^{\lambda} F(t) dt = \bar{B}_p \end{aligned}$$

from (A-5) and (A-6). Therefore, from (A-2)–(A-4) $B_p < \bar{B}_p = B - B_p$ so that $I_p(a,b) = B_p/B < 1/2$.

Theorem (A-4): If $a, b \geq 1$ then

$$1/2 < I_p(a,b) < 1 - 1/e \quad \text{when } a < b$$

$$1/e < I_p(a,b) < 1/2 \quad \text{when } a > b.$$

Proof: Immediate from (4) and theorems (A-2) and (A-3).

APPENDIX B

MONOTONIC PROPERTIES OF $I_x(a,b)$

MONOTONIC PROPERTIES OF $I_X(a,b)$

The purpose of this appendix is to show that $I_X(a,b)$ is a decreasing function of a and an increasing function of b .

Theorem B: If $a > 0$, $b > 0$, and $0 \leq x \leq 1$, then $\frac{\partial I_X(a,b)}{\partial a} \leq 0$.

Proof: Let

$$f(t;a,b) \equiv t^{a-1}(1-t)^{b-1},$$

hereafter simply denoted by $f(t)$. We have

$$I_X(a,b) \equiv B_X(a,b)/B(a,b)$$

$$B_X(a,b) \equiv \int_0^x t^{a-1}(1-t)^{b-1} dt, \quad B(a,b) \equiv \int_0^1 t^{a-1}(1-t)^{b-1} dt.$$

Then

$$\frac{\partial I_X(a,b)}{\partial a} = \frac{B(a,b) \int_0^x u^{a-1}(1-u)^{b-1} \ln u du - B_X(a,b) \int_0^1 t^{a-1}(1-t)^{b-1} \ln t dt}{B(a,b) B(a,b)}$$

$$\begin{aligned} [B(a,b)]^2 \frac{\partial I_X(a,b)}{\partial a} &= \int_0^1 \left[\int_0^x t^{a-1}(1-t)^{b-1} u^{a-1}(1-u)^{b-1} (\ln u - \ln t) du \right] dt \\ &= \int_0^x \int_0^x [f(t)f(u)(\ln u - \ln t) du] dt + \int_x^1 \int_0^x [f(t)f(u)(\ln u - \ln t) du] dt. \end{aligned}$$

But the first double integral on the right is zero. Indeed, with

$$T_{u,t} \equiv \int_0^x \int_0^x [f(t)f(u)(\ln u - \ln t) du] dt,$$

where the subscripts are used to indicate the order of integration (u then t), we have, by renaming the variables,

$$T_{u,t} = T_{t,u}.$$

Interchanging the order of integration then gives

$$T_{u,t} = T_{t,u} = -T_{u,t},$$

which implies that $T_{u,t} = 0$. Therefore,

$$[B(a,b)]^2 \frac{\partial I_X(a,b)}{\partial a} = \int_x^1 \int_0^x f(t) f(u) (\ln u - \ln t) du dt, \quad (B-1)$$

which is negative since $\ln u - \ln t \leq 0$ for $u \leq t$. Thus,

$$\frac{\partial I_X(a,b)}{\partial a} \leq 0 \text{ for all } a > 0, \quad b > 0, \quad 0 \leq x \leq 1.$$

Remarks. A more direct proof of (B-1) can be obtained by differentiating $1/[1+I_y(b,a)/I_x(a,b)]$. Also, since $\ln u - \ln t \leq u - t \leq 0$ for $0 < u \leq t \leq 1$, from (B-1) we obtain the slightly stronger inequality

$$\begin{aligned}\frac{\partial I_x(a,b)}{\partial a} &\leq \int_x^1 \int_0^x [f(t)f(u)(u-t)du]dt / [B(a,b)]^2 \\ &= \frac{B(a,b)B_x(a+1,b)}{[B(a,b)]^2} - \frac{B(a+1,b)B_x(a,b)}{[B(a,b)]^2} \\ &= \frac{a}{a+b} [I_x(a+1,b) - I_x(a,b)].\end{aligned}$$

Corollary: If $a_1 \leq a_2$ then $I_x(a_2,b) \leq I_x(a_1,b)$.

Corollary: If $b_1 \leq b_2$ then $I_x(a,b_1) \leq I_x(a,b_2)$.

Proof: Since $I_y(b,a)$ decreases as b increases, $I_x(a,b) = 1 - I_y(b,a)$ increases as b increases.

We close by noting that precisely the same technique used in Theorem B establishes a corresponding result for the incomplete gamma function. If

$$P(a,x) = \frac{1}{\Gamma(a)} \int_0^x e^{-t} t^{a-1} dt, \quad a > 0, \quad x \geq 0,$$

then $\frac{\partial P(a,x)}{\partial a} \leq 0$, and hence if $a_1 \leq a_2$ then $P(a_2,x) \leq P(a_1,x)$.

APPENDIX C

UNIFORM ASYMPTOTIC BEHAVIOR OF (16)

UNIFORM ASYMPTOTIC BEHAVIOR OF (16)

The purpose of this appendix is to show that (16) has uniform asymptotic behavior for $a \leq b$ and $a(1+a/b) \rightarrow \infty$. The proof will depend on showing that the n^{th} derivative of the function $f(v)=v/s$ ($v \geq 0$) has a bound not depending on $h=a/b$ when $a \leq b$ and n is odd.

Theorem (C-1): If $a \leq b$ then $0 \leq f'(v) < 1$ for $v \geq 0$.

Proof: Since $\frac{ds}{dv} = \frac{v}{s}(1-s)(1+hs)$ and $f = \frac{v}{s}$ we have

$$f'(v) = \frac{df}{dv} = \frac{1}{s} - \frac{v}{s^2} \frac{ds}{dv} = \frac{1}{s} - \frac{v^2}{s^3} (1-s)(1+hs). \quad (\text{C-1})$$

Also

$$\frac{v^2}{s^2} (1-s)(1+hs) = 1 - \sum_{n \geq 1} \Delta_n s^n \quad (\text{C-2})$$

$$\frac{\Delta_n}{2q} = \frac{1+h^{n+1}}{(n+1)(n+2)} + \frac{h+h^n}{n(n+1)} > 0 \quad n \text{ even} \quad (\text{C-3})$$

$$\frac{\Delta_n}{2q} = \frac{1-h^{n+1}}{(n+1)(n+2)} + \frac{h-h^n}{n(n+1)} \geq 0 \quad n \text{ odd.}$$

Therefore $f'(v) = \sum_{n \geq 1} \Delta_n s^{n-1} > 0$ is an increasing function for $s > 0$ (and hence for $v > 0$). From (C-2)

$1 - \sum_{n \geq 1} \Delta_n s^n > 0$ for $0 \leq s < 1$, so that $\sum_{n \geq 1} \Delta_n \leq 1$. Consequently, $f'(v) = \sum_{n \geq 1} \Delta_n s^{n-1} < \sum_{n \geq 1} \Delta_n \leq 1$ for $v \geq 0$.

Theorem (C-2): If $a \leq b$ then $f'(v) \rightarrow 1$ when $v \rightarrow \infty$.

Proof: From (C-3) $\sum_{n \geq 1} \Delta_n / (2q) = (1+h) \sum_{n \geq 2} 1/[n(n+1)]$, where

$$\sum_{n \geq 1} 1/[n(n+1)] = \sum_{n \geq 1} [1/n - 1/(n+1)] = 1.$$

From (16.7) we have

$$v^2 = \frac{2}{1+h} [-\ln(1-s) - \frac{1}{h} \ln(1+hs)]. \quad (\text{C-4})$$

Consequently, we now consider v as a function defined for $0 < h \leq 1$ and $0 \leq s \leq 1$.

Lemma (C-1): If $n=0, 1, 2, \dots$ then $v^n (1-s)$ can be extended to a continuous function $\phi_n(h,s)$ for $0 \leq h \leq 1$ and $0 \leq s \leq 1$. Also, $\phi_n(h,1) = 0$ for $0 \leq h \leq 1$.

Proof: From (C-4) we obtain

$$v^{2n}(1-s)^2 = \frac{2^n}{(1+h)^n} \sum_{k=0}^n \binom{n}{k} \alpha_k(s) [sg(hs)]^{n-k}$$

$$\alpha_k(s) = [-\ln(1-s)]^k (1-s)^2 \quad 0 \leq s < 1$$

$$g(z) = -\ln(1+z)/z \quad 0 < z \leq 1.$$

By L'Hopital's rule $g(z) \rightarrow -1$ when $z \rightarrow 0$. Hence, if $\bar{g}(z) = g(z)$ for $0 < z \leq 1$ and $\bar{g}(0) = -1$, then $\bar{g}(z)$ is continuous for $0 \leq z \leq 1$. Also if $w = -\ln(1-s)$ then

$$\lim_{s \rightarrow 1^-} \alpha_k(s) = \lim_{w \rightarrow \infty} w^k \exp(-2w) = 0.$$

Thus, if $\bar{\alpha}_k(s) = \alpha_k(s)$ for $0 \leq s < 1$ and $\bar{\alpha}_k(1) = 0$, then

$$\phi_n(h,s) = \left[\frac{2^n}{(1+h)^n} \sum_{k=0}^n \binom{n}{k} \bar{\alpha}_k(s) [sg(hs)]^{n-k} \right]^{1/2}$$

is a continuous extension of $v^n(1-s)$ for $0 \leq h \leq 1$ and $0 \leq s \leq 1$. Also, $\phi_n(h,1) = 0$ since each $\bar{\alpha}_k(1) = 0$.

Theorem (C-3): If $n \geq 2$ then $f^{(n)}(v)$ can be extended to a function that is continuous for $0 \leq h \leq 1$ and $0 < s \leq 1$. Also, $f^{(n)}(v) \rightarrow 0$ when $v \rightarrow \infty$.

Proof: Given a fixed value $0 < h \leq 1$. Then for any polynomial P in s (and h), $k=0,1,2,\dots$, and $m=1,2,\dots$,

$$\begin{aligned} \frac{d}{dv} \left[\frac{v^k}{s^m} (1-s) P \right] &= k \frac{v^{k-1}}{s^m} (1-s) P - m \frac{v^{k+1}}{s^{m+2}} (1-s)^2 (1+hs) P \\ &\sim \frac{v^{k+1}}{s^{m+1}} (1-s)(1+hs) P + \frac{v^{k+1}}{s^{m+1}} (1-s)^2 (1+hs) P' \end{aligned}$$

where P' is the derivative dP/ds . By induction (starting with C-1) it follows for $n \geq 2$ that $f^{(n)}(v)$ is a finite sum of terms of the form $(v^k/s^m)(1-s)P$. Consequently, by Lemma (C-1) $f^{(n)}(v)$ has a continuous extension $\psi_n(h,s)$ for $0 \leq h \leq 1$ and $0 < s \leq 1$. Also, $\psi_n(h,1) = 0$ so that $f^{(n)}(v) \rightarrow 0$ when $v \rightarrow \infty$.

Remark. For $n \geq 2$, since $f(v) = \sum_{k \geq 0} d_k v^k$ in a neighborhood of 0 and $f^{(n)}(v) \rightarrow 0$ when $v \rightarrow \infty$, by

Theorem (C-3) $f^{(n)}(v)$ is bounded for $v \geq 0$. However, the bound may depend on h . We now prove that a bound exists not depending on h when n is odd.

Definition: Given a series $\sum_{n \geq 0} \alpha_n(h)s^n$ where each $\alpha_n(h)$ is continuous for $0 \leq h \leq 1$.

Let $M_n = \max \{ |\alpha_n(h)| : 0 \leq h \leq 1 \}$. Then the series $\sum \alpha_n s^n$ is said to satisfy condition (*) when

$$\sum_{n \geq 0} M_n |s|^n \text{ converges for } |s| < 1.$$

Lemma (C-2): $f'(v) = \sum_{n \geq 1} \Delta_n s^{n-1}$ satisfies condition (*).

Proof: From (C-3) Δ_n is continuous for $0 \leq h \leq 1$ and

$$|\Delta_n| \leq \frac{4}{(n+1)(n+2)} + \frac{4}{n(n+1)}.$$

Also $\sum_{n \geq 1} 1/[n(n+1)] = 1$.

Theorem (C-4): For k odd, the series $\sum_{n \geq 0} \alpha_n(h) s^n$ for $f^{(k)}(v)$ satisfies condition (*).

Proof: By Lemma (C-2), the theorem is true for $k=1$. Assume inductively that it is true for odd k . If $e_n = \max\{ |\alpha_n(h)| : 0 \leq h \leq 1 \}$ then let $g(s) = \sum_{n \geq 0} e_n s^n$. Now

$$\begin{aligned} f^{(k+1)}(v) &= \left(\sum_{n \geq 1} n \alpha_n s^{n-1} \right) \frac{v}{s} (1-s)(1+hs) = \frac{v}{s} \sum_{n \geq 0} \beta_n(h) s^n \\ \beta_0 &= \alpha_1, \quad \beta_n = (n+1)\alpha_{n+1} - n(1-h)\alpha_n - h(n-1)\alpha_{n-1} \quad (n \geq 1). \end{aligned} \tag{C-5}$$

Since $\alpha_n(h)$ is continuous for $0 \leq h \leq 1$, therefore $\beta_n(h)$ is continuous for $0 \leq h \leq 1$. Also $|\beta_n| \leq M_n$ where $M_0 = e_1$ and $M_n = (n+1)e_{n+1} + ne_n + (n-1)e_{n-1}$ ($n \geq 1$). Since $\sum_{n \geq 0} M_n s^n$ is the series for $g'(s)(1+s+s^2)$, $\sum_{n \geq 0} M_n s^n$ converges for $|s| < 1$ and $\sum_{n \geq 0} \beta_n s^n$ satisfies condition (*).

From (C-5) we obtain

$$\begin{aligned} f^{(k+2)}(v) &= f'(v) \sum_{n \geq 0} \beta_n s^n + \sum_{n \geq 1} n \beta_n s^{n-1} \frac{v^2}{s^2} (1-s)(1+hs) \\ &= \sum_{n \geq 1} \Delta_n s^{n-1} \sum_{i \geq 0} \beta_i s^i + \sum_{n \geq 1} n \beta_n s^{n-1} \left(1 - \sum_{i \geq 1} \Delta_i s^i \right). \end{aligned}$$

Consequently, $f^{(k+2)}(v) = \sum_{n \geq 0} \delta_n(h) s^n$ where

$$\begin{aligned} \delta_0 &= \beta_1 + \beta_0 \Delta_1 \\ \delta_n &= (n+1)\beta_{n+1} + \beta_0 \Delta_{n+1} - \sum_{i=0}^{n-1} i \beta_{i+1} \Delta_{n-i} \quad (n \geq 1). \end{aligned} \tag{C-6}$$

Then δ_n is continuous for $0 \leq h \leq 1$. Also, if $c_n = \max\{ |\Delta_n| : 0 \leq h \leq 1 \}$ then

$$\begin{aligned} |\delta_n| &\leq (n+1)M_{n+1} + M_0 c_{n+1} + N_n \\ N_n &= \sum_{i=0}^{n-1} i M_{i+1} c_{n-i} \quad (n \geq 1), \quad N_0 = 0. \end{aligned}$$

Now $\sum_{n \geq 0} (n+1)M_{n+1} s^n$ converges for $|s| < 1$ since it is the derivative of $\sum_{n \geq 0} M_n s^n$,

and $\sum_{n \geq 0} c_{n+1} s^n$ converges for $|s| < 1$ by Lemma C-2. Also

$$\sum_{n \geq 0} N_n |s|^n = \sum_{n \geq 0} n M_{n+1} |s|^n \sum_{n \geq 0} c_n |s|^n \leq \sum_{n \geq 0} (n+1) M_{n+1} |s|^n \sum_{n \geq 0} c_n |s|^n,$$

which converges for $|s| < 1$. Consequently, $f^{(k+2)}(v) = \sum_{n \geq 0} \delta_n s^n$ satisfies condition (*).

Theorem (C-5): If k is odd then $f^{(k)}(v)$ can be extended to a function that is continuous for $0 \leq h \leq 1$ and $0 \leq s \leq 1$. Consequently, $|f^{(k)}(v)|$ is bounded for $v \geq 0$ by a value that does not depend on h .

Proof: It has already been verified that $f^{(k)}(v)$ can be extended to a function that is continuous for $0 \leq h \leq 1$ and $0 < s \leq 1$. Let $0 < s_0 < 1$. Since $f^{(k)}(v) = \sum_{n \geq 0} \alpha_n(h) s^n$ satisfies condition (*),

$\sum_{n \geq 0} M_n s_0^n$ converges for $M_n = \max\{|\alpha_n(h)| : 0 \leq h \leq 1\}$. Therefore, by the Weierstrass M-test,

$\sum_{n \geq 0} \alpha_n s^n$ converges uniformly for $0 \leq h \leq 1$ and $0 \leq s \leq s_0$. Since each $\alpha_n(h) s^n$ is continuous for

$0 \leq h \leq 1$ and $0 \leq s \leq s_0$, therefore $\sum_{n \geq 0} \alpha_n s^n$ is continuous for $0 \leq h \leq 1$ and $0 \leq s \leq s_0$.

APPENDIX D

FORTRAN LISTING OF BRATIO

FORTRAN LISTING OF BRATIO

In this appendix the code for BRATIO is given. The following functions are used.

$$\text{ERF}(x) = \text{erf } x$$

$$\text{ERFC1}(i,x) = \begin{cases} \text{erfc } x & \text{if } i=0 \\ \exp(x^2)\text{erfc } x & \text{if } i \neq 0 \end{cases}$$

$$\text{REXP}(x) = \exp(x) - 1$$

$$\text{ALNREL}(a) = \ln(1+a) \quad a > -1$$

$$\text{RLOG1}(a) = a - \ln(1+a) \quad a > -1$$

$$\text{GAM1}(a) = 1/\Gamma(1+a) - 1 \quad -.5 \leq a \leq 1.5$$

$$\text{GAMLN}(x) = \ln \Gamma(x) \quad x > 0$$

$$\text{GAMLN1}(a) = \ln \Gamma(1+a) \quad -.2 \leq a \leq 1.25$$

$$\text{ALGDIV}(a,b) = \ln[\Gamma(b)/\Gamma(a+b)] \quad a \geq 0, \quad b \geq 8$$

$$\text{BCORR}(a,b) = \Delta(a) + \Delta(b) - \Delta(a+b) \quad a, b \geq 8$$

$$\text{BETALN}(a,b) = \ln B(a,b) \quad a, b > 0$$

$$\text{BRCOMP}(a,b,x,y) = x^a y^b / B(a,b) \quad a, b > 0, \quad 0 < x < 1, \quad y = 1 - x$$

These functions, written by A.H.Morris, are part of the NSWC mathematics subroutine library [9].

Machine – Dependent Constants

The function SPMPAR provides the machine-dependent constants needed by BRATIO. It is necessary that SPMPAR be properly defined for the computer arithmetic being used. The constants are defined in the in-line documentation of SPMAR. Values for these constants are given in the in-line documentation. SPMPAR, released by Argonne National Laboratory, is an adaptation of the Bell Laboratories function R1MACH [D1].

Transportability

All coding adheres to the 1966 and 1977 ANSI FORTRAN standards. It is assumed that a floating point arithmetic of 6 or more digits is being used. The codes were designed specifically for k-digit arithmetics where $6 \leq k \leq 14$. If $k > 14$ then only 14-digit accuracy will normally be obtained.

Reference

- D1. Fox, P.A., Hall, A.D., and Schryer, N.L. The PORT Mathematical Subroutine Library, *ACM Trans. Math Software 4* (1978), pp. 104-126.

```

SUBROUTINE BRATIO(A, B, X, Y, W, W1, IERR)
C-----
C          EVALUATION OF THE INCOMPLETE BETA FUNCTION IX(A,B)
C
C-----  

C          IT IS ASSUMED THAT A AND B ARE NONNEGATIVE, AND THAT X .LE. 1
C          AND Y = 1 - X.  BRATIO ASSIGNS W AND W1 THE VALUES
C
C          W   = IX(A,B)
C          W1 = 1 - IX(A,B)
C
C          IERR IS A VARIABLE THAT REPORTS THE STATUS OF THE RESULTS.
C          IF NO INPUT ERRORS ARE DETECTED THEN IERR IS SET TO 0 AND
C          W AND W1 ARE COMPUTED. OTHERWISE, IF AN ERROR IS DETECTED,
C          THEN W AND W1 ARE ASSIGNED THE VALUE 0 AND IERR IS SET TO
C          ONE OF THE FOLLOWING VALUES ...
C
C          IERR = 1  IF A OR B IS NEGATIVE
C          IERR = 2  IF A = B = 0
C          IERR = 3  IF X .LT. 0 OR X .GT. 1
C          IERR = 4  IF Y .LT. 0 OR Y .GT. 1
C          IERR = 5  IF X + Y .NE. 1
C          IERR = 6  IF X = A = 0
C          IERR = 7  IF Y = B = 0
C
C-----  

C          WRITTEN BY ALFRED H. MORRIS, JR.
C          NAVAL SURFACE WEAPONS CENTER
C          DAHLGREN, VIRGINIA
C          REVISED ... JUNE 1988
C-----  

C          REAL LAMBDA
C-----  

C          ***** EPS IS A MACHINE DEPENDENT CONSTANT. EPS IS THE SMALLEST
C          FLOATING POINT NUMBER FOR WHICH 1.0 + EPS .GT. 1.0
C
C          EPS = SPMPAR(1)
C-----  

C          W = 0.0
C          W1 = 0.0
C          IF (A .LT. 0.0 .OR. B .LT. 0.0) GO TO 300
C          IF (A .EQ. 0.0 .AND. B .EQ. 0.0) GO TO 310
C          IF (X .LT. 0.0 .OR. X .GT. 1.0) GO TO 320
C          IF (Y .LT. 0.0 .OR. Y .GT. 1.0) GO TO 330
C          Z = DBLE(X) + DBLE(Y) - 1.0
C          IF (ABS(Z) .GT. EPS) GO TO 340
C
C          IERR = 0
C          IF (X .EQ. 0.0) GO TO 200
C          IF (Y .EQ. 0.0) GO TO 210
C          IF (A .EQ. 0.0) GO TO 211
C          IF (B .EQ. 0.0) GO TO 201
C
C          IND = 0
C          AO = A

```

```

BO = B
XO = X
YO = Y
EPS = AMAX1(EPS, 1.E-15)
IF (AMIN1(AO, BO) .GT. 1.0) GO TO 30
C
C          PROCEDURE FOR AO .LE. 1 OR BO .LE. 1
C
IF (X .LE. 0.5) GO TO 10
IND = 1
AO = B
BO = A
XO = Y
YO = X
C
10 IF (AMAX1(AO, BO) .GT. 1.0) GO TO 20
IF (AO .GE. AMIN1(0.2, BO)) GO TO 100
IF (XO**AO .LE. 0.9) GO TO 100
IF (XO .GE. 0.3) GO TO 110
N = 20
GO TO 130
C
20 IF (BO .LE. 1.0) GO TO 100
IF (XO .GE. 0.3) GO TO 110
IF (XO .GE. 0.1) GO TO 21
IF ((XO*BO)**AO .LE. 0.7) GO TO 100
21 IF (BO .GT. 15.0) GO TO 131
N = 20
GO TO 130
C
C          PROCEDURE FOR AO .GT. 1 AND BO .GT. 1
C
30 IF (A .GT. B) GO TO 31
LAMBDA = A - (A + B)*X
GO TO 32
31 LAMBDA = (A + B)*Y - B
32 IF (LAMBDA .GE. 0.0) GO TO 40
IND = 1
AO = B
BO = A
XO = Y
YO = X
LAMBDA = ABS(LAMBDA)
C
40 IF (BO .LT. 40.0 .AND. BO*XO .LE. 0.7) GO TO 100
IF (BO .LT. 40.0) GO TO 140
IF (AO .GT. BO) GO TO 50
IF (AO .LE. 100.0) GO TO 120
IF (LAMBDA .GT. 0.03*AO) GO TO 120
GO TO 180
50 IF (BO .LE. 100.0) GO TO 120
IF (LAMBDA .GT. 0.03*BO) GO TO 120
GO TO 180
C
C          EVALUATION OF THE APPROPRIATE ALGORITHM
C
100 W = BPSER(AO, BO, XO, EPS)

```

```

W1 = 0.5 + (0.5 - W)
GO TO 220
C
110 W1 = BPSER(BO, AO, YO, EPS)
W = 0.5 + (0.5 - W1)
GO TO 220
C
120 W = BFRAC(AO, BO, XO, YO, LAMBDA, 15.0*EPS)
W1 = 0.5 + (0.5 - W)
GO TO 220
C
130 W1 = BUP(BO, AO, YO, XO, N, EPS)
BO = BO + N
131 CALL BGRAT(BO, AO, YO, XO, W1, 15.0*EPS, IERR1)
W = 0.5 + (0.5 - W1)
GO TO 220
C
140 N = BO
BO = BO - N
IF (BO .NE. 0.0) GO TO 141
N = N - 1
BO = 1.0
141 W = BUP(BO, AO, YO, XO, N, EPS)
IF (XO .GT. 0.7) GO TO 150
W = W + BPSER(AO, BO, XO, EPS)
W1 = 0.5 + (0.5 - W)
GO TO 220
C
150 IF (AO .GT. 15.0) GO TO 151
N = 20
W = W + BUP(AO, BO, XO, YO, N, EPS)
AO = AO + N
151 CALL BGRAT(AO, BO, XO, YO, W, 15.0*EPS, IERR1)
W1 = 0.5 + (0.5 - W)
GO TO 220
C
180 W = BASYM(AO, BO, XO, YO, LAMBDA, 100.0*EPS)
W1 = 0.5 + (0.5 - W)
GO TO 220
C
C          TERMINATION OF THE PROCEDURE
C
200 IF (A .EQ. 0.0) GO TO 350
201 W = 0.0
W1 = 1.0
RETURN
C
210 IF (B .EQ. 0.0) GO TO 360
211 W = 1.0
W1 = 0.0
RETURN
C
220 IF (IND .EQ. 0) RETURN
T = W
W = W1
W1 = T
RETURN

```

```
C  
C  
C          ERROR RETURN  
C  
300 IERR = 1  
     RETURN  
310 IERR = 2  
     RETURN  
320 IERR = 3  
     RETURN  
330 IERR = 4  
     RETURN  
340 IERR = 5  
     RETURN  
350 IERR = 6  
     RETURN  
360 IERR = 7  
     RETURN  
END
```

```

REAL FUNCTION BPSER(A, B, X, EPS)
C-----  

C      POWER SERIES EXPANSION FOR EVALUATING IX(A,B) WHEN B .LE. 1  

C      OR B*X .LE. 0.7.  EPS IS THE TOLERANCE USED.  

C-----  

C      REAL N  

C
C      BPSER = 0.0
C      IF (X .EQ. 0.0) RETURN
C-----  

C      COMPUTE THE FACTOR X**A/(A*BETA(A,B))
C-----  

C      AO = AMIN1(A,B)
C      IF (AO .LT. 1.0) GO TO 10
C          Z = A*ALOG(X) - BETALN(A,B)
C          BPSER = EXP(Z)/A
C          GO TO 70
C 10 BO = AMAX1(A,B)
C      IF (BO .GE. 8.0) GO TO 60
C      IF (BO .GT. 1.0) GO TO 40
C
C      PROCEDURE FOR AO .LT. 1 AND BO .LE. 1
C
C      BPSER = X**A
C      IF (BPSER .EQ. 0.0) RETURN
C
C      APB = A + B
C      IF (APB .GT. 1.0) GO TO 20
C          Z = 1.0 + GAM1(APB)
C          GO TO 30
C 20 U = DBLE(A) + DBLE(B) - 1.0
C      Z = (1.0 + GAM1(U))/APB
C
C 30 C = (1.0 + GAM1(A))*(1.0 + GAM1(B))/Z
C      BPSER = BPSER*C*(B/APB)
C      GO TO 70
C
C      PROCEDURE FOR AO .LT. 1 AND 1 .LT. BO .LT. 8
C
C 40 U = GAMLN1(AO)
C      M = BO - 1.0
C      IF (M .LT. 1) GO TO 50
C      C = 1.0
C      DO 41 I = 1,M
C          BO = BO - 1.0
C 41      C = C*(BO/(AO + BO))
C      U = ALOG(C) + U
C
C 50 Z = A*ALOG(X) - U
C      BO = BO - 1.0
C      APB = AO + BO
C      IF (APB .GT. 1.0) GO TO 51
C          T = 1.0 + GAM1(APB)
C          GO TO 52
C 51 U = DBLE(AO) + DBLE(BO) - 1.0
C      T = (1.0 + GAM1(U))/APB
C 52 BPSER = EXP(Z)*(AO/A)*(1.0 + GAM1(BO))/T

```

```

GO TO 70
C
C          PROCEDURE FOR AO .LT. 1 AND BO .GE. 8
C
60 U = GAMLN1(AO) + ALGDIV(AO,BO)
Z = A*ALOG(X) - U
BPSER = (AO/A)*EXP(Z)
70 IF (BPSER .EQ. 0.0 .OR. A .LE. 0.1*EPS) RETURN
C----- COMPUTE THE SERIES -----
C----- SUM = 0.0
N = 0.0
C = 1.0
TOL = EPS/A
100   N = N + 1.0
      C = C*(0.5 + (0.5 - B/N))*X
      W = C/(A + N)
      SUM = SUM + W .
      IF (ABS(W) .GT. TOL) GO TO 100
      BPSER = BPSER*(1.0 + A*SUM)
      RETURN
END

```

```

      REAL FUNCTION BUP(A, B, X, Y, N, EPS)
C-----  

C----- EVALUATION OF IX(A,B) - IX(A+N,B) WHERE N IS A POSITIVE INTEGER.  

C----- EPS IS THE TOLERANCE USED.  

C-----  

      REAL L
C
      BUP = BRCOMP(A,B,X,Y)/A
      IF (N .EQ. 1 .OR. BUP .EQ. 0.0) RETURN
      NM1 = N - 1
      APB = A + B
      AP1 = A + 1.0
      SUM = 1.0
      D = 1.0
C
C       LET K BE THE INDEX OF THE MAXIMUM TERM
C
      K = 0
      IF (B .LE. 1.0) GO TO 30
      IF (Y .GT. 1.E-4) GO TO 10
      K = NM1
      GO TO 20
10   R = (B - 1.0)*X/Y - A
      IF (R .LT. 1.0) GO TO 30
      K = NM1
      IF (R .LT. FLOAT(NM1)) K = R
C
C       ADD THE INCREASING TERMS OF THE SERIES
C
20   DO 21 I = 1,K
      L = I - 1
      D = ((APB + L)/(AP1 + L))*X*D
21   SUM = SUM + D
      IF (K .EQ. NM1) GO TO 40
C
C       ADD THE REMAINING TERMS OF THE SERIES
C
30   KP1 = K + 1
      DO 31 I = KP1,NM1
      L = I - 1
      D = ((APB + L)/(AP1 + L))*X*D
      SUM = SUM + D
      IF (D .LE. EPS*SUM) GO TO 40
31   CONTINUE
C
C       TERMINATE THE PROCEDURE
C
40   BUP = BUP*SUM
      RETURN
      END

```

```

REAL FUNCTION BFRAC(A, B, X, Y, LAMBDA, EPS)
C-----CONTINUED FRACTION EXPANSION FOR IX(A,B) WHEN A,B .GT. 1.
C IT IS ASSUMED THAT LAMBDA = (A + B)*Y - B.
C-----REAL LAMBDA, N
C-----BFRAC = BRCOMP(A,B,X,Y)
IF (BFRAC .EQ. 0.0) RETURN
C
C = 1.0 + LAMBDA
CO = B/A
C1 = 1.0 + 1.0/A
YP1 = Y + 1.0
C
N = 0.0
P = 1.0
S = A + 1.0
AN = 0.0
BN = 1.0
ANP1 = 1.0
BNP1 = C/C1
R = C1/C
C
C-----CONTINUED FRACTION CALCULATION
C
10   N = N + 1.0
T = N +
W = N*(B - N)*X
E = A/S
ALPHA = (P*(P + CO)*E*E)*(W*X)
E = (1.0 + T)/(C1 + T + T)
BETA = N + W/S + E*(C + N*YP1)
P = 1.0 + T
S = S + 2.0
C
C-----UPDATE AN, BN, ANP1, AND BNP1
C
T = ALPHA*AN + BETA*ANP1
AN = ANP1
ANP1 = T
T = ALPHA*BN + BETA*BNP1
BN = BNP1
BNP1 = T
C
RO = R
R = ANP1/BNP1
IF (ABS(R - RO) .LE. EPS*R) GO TO 20
C
C-----RESCALE AN, BN, ANP1, AND BNP1
C
AN = AN/BNP1
BN = BN/BNP1
ANP1 = R
BNP1 = 1.0
GO TO 10
C

```

C TERMINATION
C
20 BFRAC = BFRAC*R
RETURN
END

```

      REAL FUNCTION BRCOMP(A, B, X, Y)
C-----  

C      EVALUATION OF X**A*Y**B/BETA(A,B)  

C-----  

C      REAL LAMBDA, LNX, LNY  

C-----  

C      CONST = 1/SQRT(2*PI)  

C-----  

C      DATA CONST/.398942280401433/  

C
C      AO = AMIN1(A,B)
C      IF (AO .GE. 8.0) GO TO 100
C
C      IF (X .GT. 0.375) GO TO 10
C          LNX = ALOG(X)
C          LNY = ALNREL(-X)
C          GO TO 20
C 10 IF (Y .GT. 0.375) GO TO 11
C          LNX = ALNREL(-Y)
C          LNY = ALOG(Y)
C          GO TO 20
C 11 LNX = ALOG(X)
C          LNY = ALOG(Y)
C
C 20 IF (AO .LT. 1.0) GO TO 30
C      Z = (A*LNX + B*LNY) - BETALN(A,B)
C      BRCOMP = EXP(Z)
C      RETURN
C-----  

C      PROCEDURE FOR A .LT. 1 OR B .LT. 1
C-----  

C 30 BO = AMAX1(A,B)
C      IF (BO .GE. 8.0) GO TO 80
C      IF (BO .GT. 1.0) GO TO 60
C
C          ALGORITHM FOR BO .LE. 1
C
C      BRCOMP = EXP(A*LNX + B*LNY)
C      IF (BRCOMP .EQ. 0.0) RETURN
C
C      APB = A + B
C      IF (APB .GT. 1.0) GO TO 40
C          Z = 1.0 + GAM1(APB)
C          GO TO 50
C 40 U = DBLE(A) + DBLE(B) - 1.0D0
C      Z = (1.0 + GAM1(U))/APB
C
C 50 C = (1.0 + GAM1(A))*(1.0 + GAM1(B))/Z
C      BRCOMP = BRCOMP*(AO*C)/(1.0 + AO/BO)
C      RETURN
C
C          ALGORITHM FOR 1 .LT. BO .LT. 8
C
C 60 U = GAMLN1(AO)
C      N = BO - 1.0
C      IF (N .LT. 1) GO TO 70
C      C = 1.0

```

```

      DO 61 I = 1,N
      BO = BO - 1.0
61    C = C*(BO/(AO + BO))
      U = ALOG(C) + U
C
70   Z = (A*LNX + B*LNY) - U
      BO = BO - 1.0
      APB = AO + BO
      IF (APB .GT. 1.0) GO TO 71
      T = 1.0 + GAM1(APB)
      GO TO 72
71   U = DBLE(AO) + DBLE(BO) - 1.0D0
      T = (1.0 + GAM1(U))/APB
72   BRCOMP = AO*EXP(Z)*(1.0 + GAM1(BO))/T
      RETURN
C
C               ALGORITHM FOR BO .GE. 8
C
80   U = GAMLN1(AO) + ALGDIV(AO,BO)
      Z = (A*LNX + B*LNY) - U
      BRCOMP = AO*EXP(Z)
      RETURN
C-----.
C-----.
C-----.
C               PROCEDURE FOR A .GE. 8 AND B .GE. 8
C-----.
100  IF (A .GT. B) GO TO 101
      H = A/B
      XO = H/(1.0 + H)
      YO = 1.0/(1.0 + H)
      LAMBDA = A - (A + B)*X
      GO TO 110
101  H = B/A
      XO = 1.0/(1.0 + H)
      YO = H/(1.0 + H)
      LAMBDA = (A + B)*Y - B
C
110  E = -LAMBDA/A
      IF (ABS(E) .GT. 0.6) GO TO 111
      U = RLOG1(E)
      GO TO 120
111  U = E - ALOG(X/XO)
C
120  E = LAMBDA/B
      IF (ABS(E) .GT. 0.6) GO TO 121
      V = RLOG1(E)
      GO TO 130
121  V = E - ALOG(Y/YO)
C
130  Z = EXP(-(A*U + B*V))
      BRCOMP = CONST*SQRT(B*XO)*Z*EXP(-BCORR(A,B))
      RETURN
      END

```

```

SUBROUTINE BGRAT(A, B, X, Y, W, EPS, IERR)
C-----  

C      ASYMPTOTIC EXPANSION FOR IX(A,B) WHEN A IS LARGER THAN B.  

C      THE RESULT OF THE EXPANSION IS ADDED TO W. IT IS ASSUMED  

C      THAT A .GE. 15 AND B .LE. 1. EPS IS THE TOLERANCE USED.  

C      IERR IS A VARIABLE THAT REPORTS THE STATUS OF THE RESULTS.  

C-----  

C      REAL J, L, LNX, NU, N2  

C      REAL C(30), D(30)  

C  

C      BM1 = (B - 0.5) - 0.5  

C      NU = A + 0.5*BM1  

C      IF (Y .GT. 0.375) GO TO 10  

C          LNX = ALNREL(-Y)  

C          GO TO 11  

C 10  LNX = ALOG(X)  

C 11  Z = -NU*LNX  

C      IF (B-Z .EQ. 0.0) GO TO 100  

C  

C          COMPUTATION OF THE EXPANSION  

C          SET R = EXP(-Z)*Z**B/GAMMA(B)  

C  

C      R = B*(1.0 + GAM1(B))*EXP(B*ALOG(Z))  

C      R = R*EXP(A*LNX)*EXP(0.5*BM1*LNX)  

C      U = ALGDIV(B,A) + B*ALOG(NU)  

C      U = R*EXP(-U)  

C      IF (U .EQ. 0.0) GO TO 100  

C      CALL GRAT1(B,Z,R,P,Q,EPS)  

C  

C      V = 0.25*(1.0/NU)**2  

C      T2 = 0.25*LNX*LNX  

C      L = W/U  

C      J = Q/R  

C      SUM = J  

C      T = 1.0  

C      CN = 1.0  

C      N2 = 0.0  

C      DO 22 N = 1,30  

C          BP2N = B + N2  

C          J = (BP2N*(BP2N + 1.0)*J + (Z + BP2N + 1.0)*T)*V  

C          N2 = N2 + 2.0  

C          T = T*T2  

C          CN = CN/(N2*(N2 + 1.0))  

C          C(N) = CN  

C          S = 0.0  

C          IF (N .EQ. 1) GO TO 21  

C              NM1 = N - 1  

C              COEF = B - N  

C              DO 20 I = 1,NM1  

C                  S = S + COEF*C(I)*D(N-I)  

C              COEF = COEF + B  

C 20      D(N) = BM1*CN + S/N  

C 21      DJ = D(N)*J  

C      SUM = SUM + DJ  

C      IF (SUM .LE. 0.0) GO TO 100  

C      IF (ABS(DJ) .LE. EPS*(SUM + L)) GO TO 30  

C 22  CONTINUE

```

```
C
C           ADD THE RESULTS TO W
C
30 IERR = 0
W = W + U*SUM
RETURN
C
C           THE EXPANSION CANNOT BE COMPUTED
C
100 IERR = 1
RETURN
END
```

```

SUBROUTINE GRAT1 (A,X,R,P,Q,EPS)
REAL J, L
C-----  

C          EVALUATION OF THE INCOMPLETE GAMMA RATIO FUNCTIONS  

C          P(A,X) AND Q(A,X)  

C  

C          IT IS ASSUMED THAT A .LE. 1. EPS IS THE TOLERANCE TO BE USED.  

C          THE INPUT ARGUMENT R HAS THE VALUE E**(-X)*X**A/GAMMA(A).  

C-----  

IF (A*X .EQ. 0.0) GO TO 130
IF (A .EQ. 0.5) GO TO 120
IF (X .LT. 1.1) GO TO 10
GO TO 50
C  

C          TAYLOR SERIES FOR P(A,X)/X**A
C  

10 AN = 3.0
C = X
SUM = X/(A + 3.0)
TOL = 0.1*EPS/(A + 1.0)
11   AN = AN + 1.0
C = -C*(X/AN)
T = C/(A + AN)
SUM = SUM + T
IF (ABS(T) .GT. TOL) GO TO 11
J = A*X*((SUM/6.0 - 0.5/(A + 2.0))*X + 1.0/(A + 1.0))
C  

Z = A*ALOG(X)
H = GAM1(A)
G = 1.0 + H
IF (X .LT. 0.25) GO TO 20
IF (A .LT. X/2.59) GO TO 40
GO TO 30
20 IF (Z .GT. -.13394) GO TO 40
C  

30 W = EXP(Z)
P = W*G*(0.5 + (0.5 - J))
Q = 0.5 + (0.5 - P)
RETURN
C  

40 L = REXP(Z)
W = 0.5 + (0.5 + L)
Q = (W*J - L)*G - H
IF (Q .LT. 0.0) GO TO 110
P = 0.5 + (0.5 - Q)
RETURN
C  

C          CONTINUED FRACTION EXPANSION
C  

50 A2NM1 = 1.0
A2N = 1.0
B2NM1 = X
B2N = X + (1.0 - A)
C = 1.0
51   A2NM1 = X*A2N + C*A2NM1
B2NM1 = X*B2N + C*B2NM1
AMO = A2NM1/B2NM1

```

```
C = C + 1.0
CMA = C - A
A2N = A2NM1 + CMA*A2N
B2N = B2NM1 + CMA*B2N
ANO = A2N/B2N
IF (ABS(ANO - AMO) .GE. EPS*ANO) GO TO 51
Q = R*ANO
P = 0.5 + (0.5 - Q)
RETURN
C
C          SPECIAL CASES
C
100 P = 0.0
Q = 1.0
RETURN
C
110 P = 1.0
Q = 0.0
RETURN
C
120 IF (X .GE. 0.25) GO TO 121
P = ERF(SQRT(X))
Q = 0.5 + (0.5 - P)
RETURN
121 Q = ERFC1(0,SQRT(X))
P = 0.5 + (0.5 - Q)
RETURN
C
130 IF (X .LE. A) GO TO 100
GO TO 110
END
```

```

REAL FUNCTION BASYM(A, B, X, Y, LAMBDA, EPS)
C-----  

C   ASYMPTOTIC EXPANSION FOR IX(A,B) FOR LARGE A AND B.  

C   LAMBDA = (A + B)*Y - B AND EPS IS THE TOLERANCE USED.  

C   IT IS ASSUMED THAT LAMBDA IS NONNEGATIVE AND THAT  

C   A AND B ARE GREATER THAN OR EQUAL TO 15.  

C-----  

REAL JO, J1, LAMBDA
REAL AO(21), BO(21), C(21), D(21)
C-----  

***** NUM IS THE MAXIMUM VALUE THAT N CAN TAKE IN THE DO LOOP  

C      ENDING AT STATEMENT 50. IT IS REQUIRED THAT NUM BE EVEN.  

C      THE ARRAYS AO, BO, C, D HAVE DIMENSION NUM + 1.  

C-----  

      DATA NUM/20/  

C-----  

C   EO = 2/SQRT(PI)
C   E1 = 2**(-3/2)
C-----  

      DATA EO/1.12837916709551/, E1/.353553390593274/
C-----  

BASYM = 0.0
IF (A .GE. B) GO TO 10
H = A/B
R0 = 1.0/(1.0 + H)
R1 = (B - A)/B
W0 = 1.0/SQRT(A*(1.0 + H))
GO TO 20
10 H = B/A
R0 = 1.0/(1.0 + H)
R1 = (B - A)/A
W0 = 1.0/SQRT(B*(1.0 + H))
C-----  

20 F = A*RLOG1(-LAMBDA/A) + B*RLOG1(LAMBDA/B)
T = EXP(-F)
IF (T .EQ. 0.0) RETURN
Z0 = SQRT(F)
Z = 0.5*(Z0/E1)
Z2 = F + F
C-----  

AO(1) = (2.0/3.0)*R1
C(1) = - 0.5*AO(1)
D(1) = - C(1)
JO = (0.5/EO)*ERFC1(1,Z0)
J1 = E1
SUM = JO + D(1)*W0*J1
C-----  

S = 1.0
H2 = H*H
HN = 1.0
W = W0
ZNM1 = Z
ZN = Z2
DO 50 N = 2, NUM, 2
HN = H2*HN
AO(N) = 2.0*RO*(1.0 + H*HN)/(N + 2.0)
NP1 = N + 1

```

```

S = S + HN
AO(NP1) = 2.0*R1*S/(N + 3.0)
C
DO 41 I = N, NP1
R = -0.5*(I + 1.0)
BO(1) = R*AO(1)
DO 31 M = 2, I
BSUM = 0.0
MM1 = M - 1
DO 30 J = 1, MM1
MMJ = M - J
BSUM = BSUM + (J*R - MMJ)*AO(J)*BO(MMJ)
30      BO(M) = R*AO(M) + BSUM/M
31      C(I) = BO(I)/(I + 1.0)
C
DSUM = 0.0
IM1 = I - 1
DO 40 J = 1, IM1
IMJ = I - J
40      DSUM = DSUM + D(IMJ)*C(J)
41      D(I) = -(DSUM + C(I))
C
JO = E1*ZNM1 + (N - 1.0)*J0
J1 = E1*ZN + N*J1
ZNM1 = Z2-ZNM1
ZN = Z2-ZN
W = WO-W
TO = D(N)*W-J0
W = WO-W
T1 = D(NP1)*W-J1
SUM = SUM + (TO + T1)
IF ((ABS(TO) + ABS(T1)) .LE. EPS*SUM) GO TO 60
50      CONTINUE
C
60 U = EXP(-BCORR(A,B))
BASYM = EO*T*U*SUM
RETURN
END

```

```
C      REAL FUNCTION REXP(X)
C      -----
C      COMPUTATION OF EXP(X) - 1
C      -----
C      DATA P1/ .914041914819518E-09/, P2/ .238082361044469E-01/
C      *     Q1/- .49999999085958E+00/, Q2/ .107141568980644E+00/
C      *     Q3/- .119041179760821E-01/, Q4/ .595130811860248E-03/
C      -----
C      IF (ABS(X) .GT. 0.15) GO TO 10
C      REXP = X*((P2*X + P1)*X + 1.0)/(((Q4*X + Q3)*X + Q2)*X
C      *           + Q1)*X + 1.0))
C      RETURN
C
C      10 W = EXP(X)
C      IF (X .GT. 0.0) GO TO 20
C      REXP = (W - 0.5) - 0.5
C      RETURN
C      20 REXP = W*(0.5 + (0.5 - 1.0/W))
C      RETURN
C      END
```

```
REAL FUNCTION ALNREL(A)
C-----  
C          EVALUATION OF THE FUNCTION LN(1 + A)  
C-----  
      DATA P1/-129418923021993E+01/, P2/.405303492862024E+00/,  
*          P3/-178874546012214E-01/  
      DATA Q1/-162752256355323E+01/, Q2/.747811014037616E+00/,  
*          Q3/-845104217945565E-01/  
C-----  
      IF (ABS(A) .GT. 0.375) GO TO 10  
      T = A/(A + 2.0)  
      T2 = T*T  
      W = (((P3*T2 + P2)*T2 + P1)*T2 + 1.0)/  
*          (((Q3*T2 + Q2)*T2 + Q1)*T2 + 1.0)  
      ALNREL = 2.0*T*W  
      RETURN  
C-----  
10   X = 1.0D0 + DBLE(A)  
      ALNREL = ALOG(X)  
      RETURN  
      END
```

```

C      REAL FUNCTION RLOG1(X)
C      -----
C      COMPUTATION OF X - LN(1 + X)
C      -----
C      DATA A/.566749439387324E-01/
C      DATA B/.456512608815524E-01/
C      -----
C      DATA P0/.333333333333333E+00/, P1/-224696413112536E+00/
C      *      P2/.620886815375787E-02/
C      DATA Q1/-127408923933623E+01/, Q2/.354508718369557E+00/
C      -----
C      IF (X .LT. -0.39 .OR. X .GT. 0.57) GO TO 100
C      IF (X .LT. -0.18) GO TO 10
C      IF (X .GT. 0.18) GO TO 20
C
C      ARGUMENT REDUCTION
C
C      H = X
C      W1 = 0.0
C      GO TO 30
C
C      10 H = DBLE(X) + 0.3D0
C      H = H/0.7
C      W1 = A - H*0.3
C      GO TO 30
C
C      20 H = 0.75D0*DBLE(X) - 0.25D0
C      W1 = B + H/3.0
C
C      SERIES EXPANSION
C
C      30 R = H/(H + 2.0)
C      T = R*R
C      W = ((P2*T + P1)*T + P0)/((Q2*T + Q1)*T + 1.0)
C      RLOG1 = 2.0*T*(1.0/(1.0 - R) - R*W) + W1
C      RETURN
C
C      100 W = (X + 0.5) + 0.5
C      RLOG1 = X - ALOG(W)
C      RETURN
C      END

```

```

REAL FUNCTION ERF (X)
C-----  

C          EVALUATION OF THE REAL ERROR FUNCTION  

C-----  

      REAL A(4),B(4),P(8),Q(8),R(5),S(5)  

C-----  

      DATA C/.564189583547756/  

C-----  

      DATA A(1)/-1.65581836870402E-4/, A(2)/3.25324098357738E-2/,  

      *     A(3)/1.02201136918406E-1/, A(4)/1.12837916709552E00/  

      DATA B(1)/4.64988945913179E-3/, B(2)/7.01333417158511E-2/,  

      *     B(3)/4.23906732683201E-1/, B(4)/1.00000000000000E00/  

      DATA P(1)/-1.36864857382717E-7/, P(2)/5.64195517478974E-1/,  

      *     P(3)/7.21175825088309E00/, P(4)/4.3162227220567E01/,  

      *     P(5)/1.52989285046940E02/, P(6)/3.39320816734344E02/,  

      *     P(7)/4.51918953711873E02/, P(8)/3.00459261020162E02/  

      DATA Q(1)/1.00000000000000E00/, Q(2)/1.27827273196294E01/,  

      *     Q(3)/7.70001529352295E01/, Q(4)/2.77585444743988E02/,  

      *     Q(5)/6.38980264465631E02/, Q(6)/9.31354094850610E02/,  

      *     Q(7)/7.90950925327898E02/, Q(8)/3.00459260956983E02/  

      DATA R(1)/2.10144126479064E00/, R(2)/2.62370141675169E01/,  

      *     R(3)/2.13688200555087E01/, R(4)/4.65807828718470E00/,  

      *     R(5)/2.82094791773523E-1/  

      DATA S(1)/9.41537750555460E01/, S(2)/1.87114811799590E02/,  

      *     S(3)/9.90191814623914E01/, S(4)/1.80124575948747E01/,  

      *     S(5)/1.00000000000000E00/  

C-----  

      AX = ABS(X)  

      IF (AX .GE. 0.5) GO TO 10  

      T = X*X  

      TOP = ((A(1)*T + A(2))*T + A(3))*T + A(4)  

      BOT = ((B(1)*T + B(2))*T + B(3))*T + B(4)  

      ERF = X*TOP/BOT  

      RETURN  

C  

10 IF (AX .GT. 4.0) GO TO 20  

      TOP = (((((P(1)*AX + P(2))*AX + P(3))*AX + P(4))*AX + P(5))*AX  

      *           + P(6))*AX + P(7))*AX + P(8)  

      BOT = (((((Q(1)*AX + Q(2))*AX + Q(3))*AX + Q(4))*AX + Q(5))*AX  

      *           + Q(6))*AX + Q(7))*AX + Q(8)  

      ERF = 0.5 + (0.5 - EXP(-X*X)*TOP/BOT)  

      IF (X .LT. 0.0) ERF = -ERF  

      RETURN  

C  

20 ERF = 1.0  

      IF (AX .GE. 5.6) GO TO 21  

      X2 = X*X  

      T = 1.0/X2  

      TOP = (((R(1)*T + R(2))*T + R(3))*T + R(4))*T + R(5)  

      BOT = (((S(1)*T + S(2))*T + S(3))*T + S(4))*T + S(5)  

      ERF = (C - TOP/(X2*BOT)) / AX  

      ERF = 0.5 + (0.5 - EXP(-X2)*ERF)  

21 IF (X .LT. 0.0) ERF = -ERF  

      RETURN  

      END

```

```

REAL FUNCTION ERFC1 (IND, X)
C-----  

C      EVALUATION OF THE COMPLEMENTARY ERROR FUNCTION  

C
C      ERFC1(IND,X) = ERFC(X)           IF IND = 0
C      ERFC1(IND,X) = EXP(X*X)*ERFC(X)  OTHERWISE
C-----  

REAL A(4),B(4),P(8),Q(8),R(5),S(5)
DOUBLE PRECISION W
C-----  

DATA C/.564189583547756/
C-----  

DATA A(1)/-1.65581836870402E-4/, A(2)/3.25324098357738E-2/,
*   A(3)/1.02201136918406E-1/, A(4)/1.12837916709552E00/
DATA B(1)/4.64988945913179E-3/, B(2)/7.01333417158511E-2/,
*   B(3)/4.23906732683201E-1/, B(4)/1.000000000000000E00/
DATA P(1)/-1.36864857382717E-7/, P(2)/5.64195517478974E-1/,
*   P(3)/7.21175825088309E00/, P(4)/4.31622272220567E01/,
*   P(5)/1.52989285046940E02/, P(6)/3.39320816734344E02/,
*   P(7)/4.51918953711873E02/, P(8)/3.00459261020162E02/
DATA Q(1)/1.000000000000000E00/, Q(2)/1.27827273196294E01/,
*   Q(3)/7.70001529352295E01/, Q(4)/2.77585444743988E02/,
*   Q(5)/6.38980264465631E02/, Q(6)/9.31354094850610E02/,
*   Q(7)/7.90950925327898E02/, Q(8)/3.00459260956983E02/
DATA R(1)/2.10144126479064E00/, R(2)/2.62370141675169E01/,
*   R(3)/2.13688200555087E01/, R(4)/4.65807828718470E00/,
*   R(5)/2.82094791773523E-1/
DATA S(1)/9.41537750555460E01/, S(2)/1.87114811799590E02/,
*   S(3)/9.90191814623914E01/, S(4)/1.80124575948747E01/,
*   S(5)/1.000000000000000E00/
C-----  

C      ABS(X) .LT. 0.47
C
C      AX = ABS(X)
IF (AX .GE. 0.47) GO TO 10
T = X*X
TOP = ((A(1)*T + A(2))*T + A(3))*T + A(4)
BOT = ((B(1)*T + B(2))*T + B(3))*T + B(4)
ERFC1 = 0.5 + (0.5 - X*TOP/BOT)
IF (IND .NE. 0) ERFC1 = EXP(T) * ERFC1
RETURN
C
C      0.47 .LE. ABS(X) .LE. 4
C
10 IF (AX .GT. 4.0) GO TO 30
TOP = (((((P(1)*AX + P(2))*AX + P(3))*AX + P(4))*AX + P(5))*AX
*          + P(6))*AX + P(7))*AX + P(8)
BOT = (((((Q(1)*AX + Q(2))*AX + Q(3))*AX + Q(4))*AX + Q(5))*AX
*          + Q(6))*AX + Q(7))*AX + Q(8)
ERFC1 = TOP/BOT
C
20 IF (IND .EQ. 0.0) GO TO 21
IF (X .LT. 0.0) ERFC1 = 2.0*EXP(X*X) - ERFC1
RETURN
21 W = DBLE(X)*DBLE(X)
T = W

```

```

E = W - DBLE(T)
ERFC1 = ((0.5 + (0.5 - E)) * EXP(-T)) * ERFC1
IF (X .LT. 0.0) ERFC1 = 2.0 - ERFC1
RETURN
C
C           ABS(X) .GT. 4
C
30 IF (X .LE. -5.5) GO TO 40
IF (IND .EQ. 0.0 .AND. X .GT. 50.0) GO TO 50
T = (1.0/X)**2
TOP = (((R(1)*T + R(2))*T + R(3))*T + R(4))*T + R(5)
BOT = (((S(1)*T + S(2))*T + S(3))*T + S(4))*T + S(5)
ERFC1 = (C - T*TOP/BOT) / AX
GO TO 20
C
C           LIMIT VALUE FOR LARGE NEGATIVE X
C
40 ERFC1 = 2.0
IF (IND .NE. 0) ERFC1 = 2.0*EXP(X*X)
RETURN
C
C           LIMIT VALUE FOR LARGE POSITIVE X
C           WHEN IND = 0
C
50 ERFC1 = 0.0
RETURN
END

```

```

C      REAL FUNCTION GAM1(A)
C      -----
C      COMPUTATION OF 1/GAMMA(A+1) - 1 FOR -0.5 .LE. A .LE. 1.5
C      -----
C      REAL P(7), Q(5), R(9)
C      -----
C      DATA P(1)/ .577215664901533E+00/, P(2)/-.409078193005776E+00/
C      *     P(3)/-.230975380857675E+00/, P(4)/ .597275330452234E-01/
C      *     P(5)/ .766968181649490E-02/, P(6)/-.514889771323592E-02/
C      *     P(7)/ .589597428611429E-03/
C      -----
C      DATA Q(1)/ .100000000000000E+01/, Q(2)/ .427569613095214E+00/
C      *     Q(3)/ .158451672430138E+00/, Q(4)/ .261132021441447E-01/
C      *     Q(5)/ .423244297896961E-02/
C      -----
C      DATA R(1)/-.422784335098468E+00/, R(2)/-.771330383816272E+00/
C      *     R(3)/-.24475776522226E+00/, R(4)/ .118378989872749E+00/
C      *     R(5)/ .930357293360349E-03/, R(6)/-.118290993445146E-01/
C      *     R(7)/ .223047661158249E-02/, R(8)/ .266505979058923E-03/
C      *     R(9)/-.132674909766242E-03/
C      -----
C      DATA S1 / .273076135303957E+00/, S2 / .559398236957378E-01/
C      -----
C      T = A
C      D = A - 0.5
C      IF (D .GT. 0.0) T = D - 0.5
C      IF (T) 30,10,20
C
10 GAM1 = 0.0
      RETURN
C
20 TOP = (((((P(7)*T + P(6))*T + P(5))*T + P(4))*T + P(3))*T
      *           + P(2))*T + P(1)
BOT = (((Q(5)*T + Q(4))*T + Q(3))*T + Q(2))*T + 1.0
W = TOP/BOT
IF (D .GT. 0.0) GO TO 21
      GAM1 = A*W
      RETURN
21 GAM1 = (T/A)*((W - 0.5) - 0.5)
      RETURN
C
30 TOP = ((((((R(9)*T + R(8))*T + R(7))*T + R(6))*T + R(5))*T
      *           + R(4))*T + R(3))*T + R(2))*T + R(1)
BOT = (S2*T + S1)*T + 1.0
W = TOP/BOT
IF (D .GT. 0.0) GO TO 31
      GAM1 = A*(W + 0.5) + 0.5
      RETURN
31 GAM1 = T*W/A
      RETURN
      END

```

```

REAL FUNCTION GAMLN(A)
*****
C EVALUATION OF LN(GAMMA(A)) FOR POSITIVE A
*****
C WRITTEN BY ALFRED H. MORRIS
C      NAVAL SURFACE WEAPONS CENTER
C      DAHLGREN, VIRGINIA
C -----
C D = 0.5*(LN(2*PI) - 1)
C -----
C DATA D/.418938533204673/
C -----
C DATA CO/.8333333333333E-01/, C1/-27777777770481E-02/, .
1   C2/.793650663183693E-03/, C3/-595156336428591E-03/, .
2   C4/ 820756370353826E-03/
C -----
C IF (A .GT. 0.8) GO TO 10
C      GAMLN = GAMLN1(A) - ALOG(A)
C      RETURN
10 IF (A .GT. 2.25) GO TO 20
C      X = DRLE(A) - 1.0
C      GAMLN = GAMLN1(X)
C      RETURN
C
20 IF (A .GE. 15.0) GO TO 30
C      N = A - 1.25
C      X = A
C      W = 1.0
C      DO 21 I = 1,N
C          X = X - 1.0
21   W = X*W
C      GAMLN = GAMLN1(X - 1.0) + ALOG(W)
C      RETURN
C
30 X = (1.0/A)**2
C      W = (((C4*X + C3)*X + C2)*X + C1)*X + CO/A
C      GAMLN = (D + W) + (A - 0.5)*(ALOG(A) - 1.0)
C      END

```

```

C      REAL FUNCTION GAMLN1(A)
C      -----
C      EVALUATION OF LN(GAMMA(1 + A)) FOR -0.2 .LE. A .LE. 1.25
C      -----
C      DATA P0/.577215664901533E+00/, P1/.844203922187225E+00/
C      *     P2/-168860593646662E+00/, P3/-780427615533591E+00/
C      *     P4/-402055799310489E+00/, P5/-673562214325671E-01/
C      *     P6/-271935708322958E-02/
C      DATA Q1/.288743195473681E+01/, Q2/.312755088914843E+01/
C      *     Q3/.156875193295039E+01/, Q4/.361951990101499E+00/
C      *     Q5/.325038868253937E-01/, Q6/.667465618796164E-03/
C      -----
C      DATA R0/.422784335098467E+00/, R1/.848044614534529E+00/
C      *     R2/.565221050691933E+00/, R3/.156513060486551E+00/
C      *     R4/.170502484022650E-01/, R5/.497958207639485E-03/
C      DATA S1/.124313399877507E+01/, S2/.548042109832463E+00/
C      *     S3/.101552187439830E+00/, S4/.713309612391000E-02/
C      *     S5/.116165475989616E-03/
C      -----
C      IF (A .GE. 0.6) GO TO 10
C      W = (((((P6*A + P5)*A + P4)*A + P3)*A + P2)*A + P1)*A + P0)/
C      *     (((((Q6*A + Q5)*A + Q4)*A + Q3)*A + Q2)*A + Q1)*A + 1.0)
C      GAMLN1 = -A*W
C      RETURN
C
C      10 X = DBLE(A) - 1.0
C      W = (((((R5*X + R4)*X + R3)*X + R2)*X + R1)*X + R0)/
C      *     (((((S5*X + S4)*X + S3)*X + S2)*X + S1)*X + 1.0)
C      GAMLN1 = X*W
C      RETURN
C      END

```

```

REAL FUNCTION BETALN(AO, BO)
C-----  

C----- EVALUATION OF THE LOGARITHM OF THE BETA FUNCTION  

C-----  

C----- E = 0.5*LN(2*PI)  

C-----  

C----- DATA E / .918938533204673/  

C-----  

C----- A = AMIN1(AO,BO)  

C----- B = AMAX1(AO,BO)  

C----- IF (A .GE. 8.0) GO TO 60  

C----- IF (A .GE. 1.0) GO TO 20  

C-----  

C----- PROCEDURE WHEN A .LT. 1  

C-----  

C----- IF (B .GE. 8.0) GO TO 10  

C----- BETALN = GAMLN(A) + (GAMLN(B) - GAMLN(A + B))  

C----- RETURN  

C----- 10 BETALN = GAMLN(A) + ALGDIV(A,B)  

C----- RETURN  

C-----  

C----- PROCEDURE WHEN 1 .LE. A .LT. 8  

C-----  

C----- 20 IF (A .GT. 2.0) GO TO 30  

C----- IF (B .GT. 2.0) GO TO 21  

C----- BETALN = GAMLN(A) + GAMLN(B) - GSUMLN(A,B)  

C----- RETURN  

C----- 21 W = 0.0  

C----- IF (B .LT. 8.0) GO TO 40  

C----- BETALN = GAMLN(A) + ALGDIV(A,B)  

C----- RETURN  

C-----  

C----- REDUCTION OF A WHEN A .GT. 2  

C-----  

C----- 30 N = A - 1.0  

C----- W = 0.0  

C----- DO 31 I = 1,N  

C-----   A = A - 1.0  

C-----   H = A/B  

C-----   W = W + ALOG(H/(1.0 + H))  

C----- 31 CONTINUE  

C----- IF (B .LT. 8.0) GO TO 40  

C----- BETALN = W + GAMLN(A) + ALGDIV(A,B)  

C----- RETURN  

C-----  

C----- REDUCTION OF B WHEN B .LT. 8  

C-----  

C----- 40 N = B - 1.0  

C----- Z = 1.0  

C----- DO 41 I = 1,N  

C-----   B = B - 1.0  

C-----   41 Z = Z*(B/(A + B))  

C----- BETALN = W + ALOG(Z) + (GAMLN(A) + (GAMLN(B) - GSUMLN(A,B)))  

C----- RETURN  

C-----  

C----- PROCEDURE WHEN A .GE. 8

```

```
60 W = BCORR(A,B)
    H = A/B
    C = H/(1.0 + H)
    U = -(A - 0.5)*ALOG(C)
    V = B*ALNREL(H)
    IF (U .LE. V) GO TO 61
        BETALN = (((-0.5*ALOG(B) + E) + W) - V) - U
        RETURN
61 BETALN = (((-0.5*ALOG(B) + E) + W) - U) - V
    RETURN
    END
```

```
C      REAL FUNCTION GSUMLN(A,B)
C-----+
C-----+ EVALUATION OF THE FUNCTION LN(GAMMA(A + B))
C-----+ FOR 1 .LE. A .LE. 2 AND 1 .LE. B .LE. 2
C-----+
C-----+
C      X = DBLE(A) + DBLE(B) - 2.D0
C      IF (X .GT. 0.25) GO TO 10
C          GSUMLN = GAMLN1(1.0 + X)
C          RETURN
C 10 IF (X .GT. 1.25) GO TO 20
C          GSUMLN = GAMLN1(X) + ALNREL(X)
C          RETURN
C 20 GSUMLN = GAMLN1(X - 1.0) + ALOG(X*(1.0 + X))
C          RETURN
C
```

```

REAL FUNCTION BCORR(AO, BO)
-----
C EVALUATION OF DEL(AO) + DEL(BO) - DEL(AO + BO) WHERE
C LN(GAMMA(A)) = (A - 0.5)*LN(A) - A + 0.5*LN(2*PI) + DEL(A).
C IT IS ASSUMED THAT AO .GE. 8 AND BO .GE. 8.
C -----
C DATA CO/.833333333333E-01/, C1/- .277777777750991E-02/,
*      C2/.793650666825390E-03/, C3/- 595202931351870E-03/,
*      C4/.837308034031215E-03/, C5/- .165322962780713E-02/
C -----
C A = AMIN1(AO, BO)
C B = AMAX1(AO, BO)
C
C H = A/B
C C = H/(1.0 + H)
C X = 1.0/(1.0 + H)
C X2 = X*X
C
C SET SN = (1 - X**N)/(1 - X)
C
C S3 = 1.0 + (X + X2)
C S5 = 1.0 + (X + X2*S3)
C S7 = 1.0 + (X + X2*S5)
C S9 = 1.0 + (X + X2*S7)
C S11 = 1.0 + (X + X2*S9)
C
C SET W = DEL(B) - DEL(A + B)
C
C T = (1.0/B)**2
C W = (((((C5*S11*T + C4*S9)*T + C3*S7)*T + C2*S5)*T + C1*S3)*T + CO
C W = W*(C/B)
C
C COMPUTE DEL(A) + W
C
C T = (1.0/A)**2
BCORR = (((((C5*T + C4)*T + C3)*T + C2)*T + C1)*T + CO)/A + W
RETURN
END

```

```

REAL FUNCTION ALGDIV(A,B)
C-----  

C COMPUTATION OF LN(GAMMA(B))/GAMMA(A+B)) WHEN B .GE. 8  

C  

C IN THIS CODE DEL(Z) IS THE FUNCTION WHERE  

C LN(GAMMA(Z)) = (Z - 0.5)*LN(Z) - Z + 0.5*LN(2*PI) + DEL(Z)  

C-----  

C DATA CO/.83333333333333E-01/, C1/- .27777777760991E-02/,  

* C2/.793650666825390E-03/, C3/- .595202931351870E-03/,  

* C4/.837308034031215E-03/, C5/- .165322962780713E-02/  

C-----  

C IF (A .LE. B) GO TO 10
H = B/A
C = 1.0/(1.0 + H)
X = H/(1.0 + H)
D = A + (B - 0.5)
GO TO 20
10 H = A/B
C = H/(1.0 + H)
X = 1.0/(1.0 + H)
D = B + (A - 0.5)
C  

SET SN = (1 - X**N)/(1 - X)
C  

20 X2 = X*X
S3 = 1.0 + (X + X2)
S5 = 1.0 + (X + X2*S3)
S7 = 1.0 + (X + X2*S5)
S9 = 1.0 + (X + X2*S7)
S11 = 1.0 + (X + X2*S9)
C  

SET W = DEL(B) - DEL(A + B)
C  

T = (1.0/B)**2
W = (((((C5*S11*T + C4*S9)*T + C3*S7)*T + C2*S5)*T + C1*S3)*T + CO
W = W*(C/B)
C  

COMBINE THE RESULTS
C  

U = D*ALNREL(A/B)
V = A*(ALOG(B) - 1.0)
IF (U .LE. V) GO TO 30
ALGDIV = (W - V) - U
RETURN
30 ALGDIV = (W - U) - V
RETURN
END

```



```

C      MACHINE CONSTANTS FOR THE CDC 6000/7000 SERIES.
C      (INTEGER FORMAT FOR FORTRAN 4 AND 5 COMPILERS)
C
C      DATA MCHEPS(1) / 261630990852554752 /
C      DATA MINMAG(1) / 422212465065984 /
C      DATA MAXMAG(1) / 576179277326712831 /
C
C      MACHINE CONSTANTS FOR THE CRAY-1.
C
C      DATA RMACH(1) / 0377224000000000000000000B /
C      DATA RMACH(2) / 0200034000000000000000000B /
C      DATA RMACH(3) / 0577777777777777777776B /
C
C      MACHINE CONSTANTS FOR THE DATA GENERAL ECLIPSE S/200.
C
C      NOTE - IT MAY BE APPROPRIATE TO INCLUDE THE FOLLOWING CARD -
C      STATIC RMACH(3)
C
C      DATA MINMAG/20K,O/,MAXMAG/77777K,177777K/
C      DATA MCHEPS/36020K,O/
C
C      MACHINE CONSTANTS FOR THE HARRIS 220.
C
C      DATA MCHEPS(1) / '20000000, '00000353 /
C      DATA MINMAG(1) / '20000000, '00000201 /
C      DATA MAXMAG(1) / '37777777, '00000177 /
C
C      MACHINE CONSTANTS FOR THE HONEYWELL 600/6000 SERIES.
C
C      DATA RMACH(1) / 0716400000000 /
C      DATA RMACH(2) / 0402400000000 /
C      DATA RMACH(3) / 0376777777777 /
C
C      MACHINE CONSTANTS FOR THE HP 2100
C
C      DATA MCHEPS(1), MCHEPS(2) / 40000B,      327B /
C      DATA MINMAG(1), MINMAG(2) / 40000B,      1 /
C      DATA MAXMAG(1), MAXMAG(2) / 77777B, 177776B /
C
C      MACHINE CONSTANTS FOR THE HP 9000
C
C      DATA RMACH(1) / .1192093E-06 /
C      DATA RMACH(2) / .5877472E-38 /
C      DATA RMACH(3) / .3402823E+39 /
C
C      MACHINE CONSTANTS FOR THE IBM 360/370 SERIES,
C      THE AMDAHL 470/V6, THE ICL 2900, THE ITEL AS/6,
C      THE XEROX SIGMA 5/7/9 AND THE SEL SYSTEMS 85/86.
C
C      DATA RMACH(1) / Z3C100000 /
C      DATA RMACH(2) / Z00100000 /
C      DATA RMACH(3) / Z7FFFFFF /
C
C      MACHINE CONSTANTS FOR THE IBM PC - MICROSOFT FORTRAN
C
C      DATA MCHEPS(1) / #34000000 /
C      DATA MINMAG(1) / #00800000 /

```

```

C      DATA MAXMAG(1) / #7F7FFFFF /
C
C      MACHINE CONSTANTS FOR THE IBM PC ~ PROFESSIONAL FORTRAN,
C      LAHEY FORTRAN, AND RM FORTRAN
C
C      DATA MCHEPS(1) / Z'34000000' /
C      DATA MINMAG(1) / Z'00800000' /
C      DATA MAXMAG(1) / Z'7F7FFFFF' /
C
C      MACHINE CONSTANTS FOR THE PDP-10 (KA OR KI PROCESSOR).
C
C      DATA RMACH(1) / "147400000000 /
C      DATA RMACH(2) / "000400000000 /
C      DATA RMACH(3) / "377777777777 /
C
C      MACHINE CONSTANTS FOR THE PDP-11 FORTRAN SUPPORTING
C      32-BIT INTEGERS (EXPRESSED IN INTEGER AND OCTAL).
C
C      DATA MCHEPS(1) / 889192448 /
C      DATA MINMAG(1) / 8388608 /
C      DATA MAXMAG(1) / 2147483647 /
C
C      DATA RMACH(1) / 006500000000 /
C      DATA RMACH(2) / 000040000000 /
C      DATA RMACH(3) / 017777777777 /
C
C      MACHINE CONSTANTS FOR THE PDP-11 FORTRAN SUPPORTING
C      16-BIT INTEGERS (EXPRESSED IN INTEGER AND OCTAL).
C
C      DATA MCHEPS(1),MCHEPS(2) / 13568,      0 /
C      DATA MINMAG(1),MINMAG(2) / 128,       0 /
C      DATA MAXMAG(1),MAXMAG(2) / 32767,     -1 /
C
C      DATA MCHEPS(1),MCHEPS(2) / 0032400, 0000000 /
C      DATA MINMAG(1),MINMAG(2) / 0000200, 0000000 /
C      DATA MAXMAG(1),MAXMAG(2) / 0077777, 0177777 /
C
C      MACHINE CONSTANTS FOR THE UNIVAC 1100 SERIES.
C
C      DATA RMACH(1) / 0147400000000 /
C      DATA RMACH(2) / 0000400000000 /
C      DATA RMACH(3) / 0377777777777 /
C
C      MACHINE CONSTANTS FOR THE VAX 11/780
C      (EXPRESSED IN INTEGER AND HEXADECIMAL)
C
C      DATA MCHEPS(1) /      13568 /
C      DATA MINMAG(1) /      128 /
C      DATA MAXMAG(1) /     -32769 /
C
C      DATA MCHEPS(1) / Z00003500 /
C      DATA MINMAG(1) / Z00000080 /
C      DATA MAXMAG(1) / ZFFFF7FFF /
C
C      SPMPAR = RMACH(I)
C      RETURN
C

```

C LAST CARD OF FUNCTION SPMPAR.
C
END

DISTRIBUTION

<u>Copies</u>	<u>Copies</u>	
Chief of Naval Operations Attn: OP-961 OP-982 OP-982E OP-983 OP-987 Washington D.C. 20350	The Library of Congress 1 Attn: Exchange and Gift Division 1 Washington D.C. 20540 Director 1 Office of Naval Research Branch Office 1030 East Green Street Pasadena, Ca 91101	1
Commander, Naval Facilities Engineering Command Department of the Navy Attn: Code 032F 200 Stovall Street Alexandria, VA 22332	Commanding Officer 1 Marine Aviation Detachment Pacific Missile Test Center Point Mugu, CA 93042	1
Office of Naval Research Code 411 800 North Quincy Street Arlington VA 22217	Director 1 U.S. Army Ballistic Research Lab. Attn: DRDAR-TSB-S (STINFO) Aberdeen Proving Ground, MD 21005	1
Commander, Naval Sea Systems Command Attn: SEA 05R Department of the Navy Washington, DC 20362	Director 1 National Security Agency Attn: Library Fort George G. Meade, MD 20755	1
Fleet Analysis Center Naval Weapons Station Seal Beach Attn: Library Corona, CA 91720	Commanding General White Sands Missile Range Attn: Technical Library, Documents Section 1 White Sands Missile Range, NM 88002	1
Commander, U.S. Naval Weapons Center Attn: Code 38 China Lake, CA 93555	Director 1 National Bureau of Standards Attn: Library Gaithersburg, MD 20760	1
Argonne National Laboratory Attn: Dr. A. H. Jaffey Bldg. 200 9700 South Cass Avenue Argonne, IL 60439	Rand Corporation 1 Attn: Library D 1700 Main Street Santa Monica, CA 90406	1
U.S. Naval Observatory 34th Street and Mass. Ave, N.W. Attn: Library Washington, DC 20390	Columbia University Prof. H. Goldstein 1 520 W. 120th Street, Engter Bldg. New York, NY 10027	1

University of Chicago Attn: Prof. W. Kruskal Statistics Department 5734 Univ. Ave. Chicago, IL 60637	1	University of Wisconsin-Madison Prof. J. Gurland Department of Statistics 1210 West Dayton Street Madison, WI 53706	1
Joint Strategic Target Planning Staff Offut Air Force Base, NB 68113	1	Chief of Missile and Projectile Systems Design John A. Simpson Norden Systems Inc. P.O. Box 5300	1
Massachusetts Institute of Technology Attn: Computation Center Cambridge, MA 02139	1	Norwalk, CT 06856	
President Naval War College Newport, RI 02840	1	Shering Plough Corp. M. Miller 60 Orange Street Bloomfield, NJ 07003	1
University of New Orleans Dr. George Ioup, Physics Dept. Lake Front, New Orleans LA 70148	1	Department of Mathematics Prof. Joo Koo Northeastern Illinois University Chicago, IL 60625	1
Willian C. Guenther Box 3332 University Station Laramie, WY 82071	1	Trinity College Harvey S. Picker Physics Department Hartford, CT 06106	1
Sandia National Laboratories Dr. Donald E. Amos, 5642 P.O. Box 5800 Albuquerque, NM 87185	1	Naval Ocean Systems Center Attn: Library San Diego, CA 92152	1
Exploratory Technology Donald Baker Moore P.O. Box KK Fairfield, CA 94533	1	U.S. Naval Oceanographic Office Navy Lib./NSTL Station Bay St. Louis, MS 39522	1
The Johns Hopkins University Dt. I. Sugai, F4A Applied Physics Laboratory Johns Hopkins Road Laurel, MD 20707	1	General Electric Company H. Saunders Building 41, Room 304 One River Road Schenectady, NY 12345	1
Washington State University Prof. George Marsaglia Computer Science Dept. Pullman, WA 99164	1	Lockheed Missile & Space Company R. F. Hausman Department 6212-- Bldg. 104 P.O. Box 504 Sunnyvale, CA 94088	1

Dr. Harold Crutcher 35 Westall Avenue Asheville, NC 28804	1	Office of Naval Research Attn: Math & Sciences Division Library Washington, DC 20360	1
Deputy Director Marine Corps Operations Analysis Group Attn: Dr. B. Barfoot 2000 N. Beauregard Street Alexandria, VA 22311	1	Commander, David W. Taylor Naval Ship Research & Develop. Center Code 1120 (E. O'Neill) Code 1129 (E. Wolfe) Code 1127 (S. Baylin) Code 185 Code 1805 Code 5220 (Technical Library) Bethesda, MD 20084	1 1 1 1 1 1 1
LMSC Dr. H. Weingarten 0/8449 B 538 P.O. Box 504 Sunnyvale, CA 94088-3504	1	AVARDA George Cyrcos DAVAA-N-SA Fort Monmouth, NJ 07703	1
Harvard University D.C. Hoaglin Dept. of Statistics 1 Oxford Street Cambridge Mass 02138	1	U.S. Army Tradoc Systems Analysis Activity Peter Shugart Attn: ATAA-DDD White Sands Missile Range, NM 88002	1
Dr. A.V. Hershey 722 Redwood Lane Pacific Grove, CA 93950	1	Director Naval Research Laboratory Washington, D.C. 20375	1
Naval Research Laboratory Alvin B. Owens Code 2303 Washington, D.C. 20375	1		

Internal Distribution

D	1	K104D	20
D1	1	K106 (Dr. M. Thomas)	2
D2 (D. Snyder)	1	K106 (Dr. J. Crigler)	1
E	1	K106 (G. Gemmill)	1
E211	1	K107 (R. Mulliken)	1
E231	10	K33 (A. Morris)	50
F41 (E. Morgan)	1	K40	1
F411	1	K43 (R. Coppola)	1
F42 (E. Spooner)	1	N	1
G	1	N302 (S. Vittoria)	1
G102 (G. Hornbaker)	1	R44	1
K	1	R44 (Dr. A.H. Van Tuyl)	1
K10	1		